92

1893.7

# 

THIRTY-EIGHTH ANNUAL REPORT

UPON THE

## Births, Marriages and Deaths

IN THE

### CITY OF PROVIDENCE

FOR THE YEAR

1892.

BY CHARLES V. CHAPIN, M. D.,

CITY REGISTRAR.



THE LIBRARY OF THE NOV 22 1968 UNIVERSITY OF ILLINOIS

J. A. & R. A. REID, CITY PRINTERS, Dyer and Pine Streets.

1893.

wx o

## SHT SASISO SHOL YSASSLI



The series of reports on births, marriages and deaths in

3 the notes on the causes of death for the five years, 1916 to 1920, the usual tables of births covering the same period, as well general summary of births, marriages and deaths. and.

and in typography, they do render available material similar to that Although these two reports are unsatisfactory in arrangement in the earlier registration reports and which it is difficult to obtain for so long a period for other American cities The series of reports on births, marriages and deaths in Providence was begun in 1855 and continued for sixty-one years through 1915, During the next four years it was impossible to print any report. The ordinance requires the report for any year to be in print by April 1 of the succeeding year unless special authority is granted and this could not be obtained. An antiquated registration law, which happily has now been brought up to date, precluded the early tabulation of births and the inability of the printer to obtain adequate help under war conditions helped to make the publication of a report impossible.

In 1919 the printing of a report was commenced, but it could not be completed until 1920 and is designated as the Sixty-second to Sixty-sixth Report. It is very meagre but contains the fundamental tables of deaths by cause, sex and age and of deaths by use and season, for each of the five years, 1916 to 1920, as well as several other important tables of deaths for these years. Most of the tables of marriages for the five years were included, as well as the tables of infant mortality. The very useful tables called "Notes on Causes of Death," which had been printed for so many years, had to be omitted.

The present report, nominally for the year 1921, contains also the notes on the causes of death for the five years, 1916 to 1920, and the usual tables of births covering the same period, as well as the general summary of births, marriages and deaths.

Although these two reports are unsatisfactory in arrangement and in typography, they do render available material similar to that in the earlier registration reports and which it is difficult to obtain for so long a period for other American cities.

## CONTENTS.

	PAGE.
BIRTHS Wards, months, sex and color, (Table I.)	5-23
Wards, months, sex and color, (Table I.) Births in connection with season, and sex. Number and proportion of the sexes, 39 years, (Table II.) Color:—Births, marriages and deaths in 37 years. Parentage of children born in each ward, (Table III.) Number and percentage of American and of foreign children, 37 years. Parentage:—Mixture of different nations, (Table IIV.) Number of first, second. etc., children in 1892, and 1855-1892, (Table V.). Average number of children to each mother, 38 years, (Table VI.) Number of the child, with age of the mother, (Table VII.) Plurality births, statistics for 1892. Plurality births, 38 years' summary. Illegitimate children.	7-8
Number and proportion of the seves 39 years (Table II)	8
Color:—Births, marriages and deaths in 37 years.	8-9
Parentage of children born in each ward, (Table III.)	10
Number and percentage of American and of foreign children, 37 years.	11-12
Parentage: -Mixture of different nations, (Table IV.)	13
Number of first, second, etc., children in 1892, and 1855-1892, (Table V.)	14
Average number of children to each mother, 38 years, (Table VI.)	15
Number of the chick, with age of the mother, (Table VII.)	16-17
Plurality Dirtils, Statistics for 1092.	19-20 20-23
Illegitime to shidren	23
megitimate children	40
MARRIAGES	24-34
MARRIAGES.  Marriages at different seasons, 38 years, (Table VIII.).  Birth-places of parties married in 1892, (Table IX.).  Bride and groom born in the same country, etc.  Proportions of American, foreign and mixed marriages, 38 years.  Marriages of colored persons	24
Birth-places of parties married in 1892. (Table IX.)	26
Bride and groom born in the same country, etc	26-27
Proportions of American, foreign and mixed marriages, 38 years	26-27 27
Marriages of colored persons	28
Age of persons married in 1892 (Table X.)	29
Average age of persons married, 37 years, (Table)	30
Number of times married, (Table XI.)	30
Marriages of divorced persons	31
Netwitz and residence	32
Proportions of American, foreign and mixed marriages, 38 years.  Marriages of colored persons.  Age of persons married in 1892 (Table X.).  Average age of persons married, 37 years, (Table).  Number of times married, (Table XI.).  Marriages of divorced persons.  Denominational  Nativity and residence.  Marriage and education; certificate signed with a mark.	33-34
Mairiage and education, certificate signed with a mark	00-01
DEATHS	25_102
Deaths to population in 1892, proportion in each class	35
Deaths by sex condition, color, nativity parentage, and locality	. 00
(Table XII.)	36
Deaths by season: numbers and proportions: recapitulation	37
Sex, condition, color, nativity, parentage, and locality of decedents, 189	38-40
Aggregate and average age by parentage, 1892, (Table XIII.)	41
Aggregate and average age by parentage, 37 years, (Table XIV.)	42
Percentages at different ages by parentage, 1892, (Table XV.)	43
Percentages under live years of age by parentage, 37 years, (Table XVI.	45
Percentages at different ages by parentage, at years, (Table A VII.)	47
YVIII)	48
Deaths to population in 1892, proportion in each class.  Deaths by sex, condition, color, nativity, parentage, and locality (Table XII.).  Deaths by season; numbers and proportions; recapitulation  Sex, condition, color, nativity, parentage, and locality of decedents, 189  Aggregate and average age by parentage, 1892, (Table XIII.)  Aggregate and average age by parentage, 1892, (Table XIV.)  Percentages at different ages by parentage, 1892, (Table XV.)  Percentages under five years of age by parentage, 37 years, (Table XVI.)  Deaths by age and parentage in each month and quarter, 1892, (Table XVIII.)  Causes of death, sex, age, parentage and percentages, (Table XIX.).	. 50-54
Causes of death, sex, age, parentage and percentages, (Table XIX.) Causes of death in connection with season, (Table XX.)	55-56
Notes on the Causes of Death in 1892.  Bronchitis; statistics for 37 years, (Table).  Cancer; statistics of sex and parentage, 37 years.  Cancer that they dead out to provide the parentage of t	57-102
Bronchitis; statistics for 37 years, (Table)	61-62
Cancer; statistics of sex and parentage, 37 years	63
Cnolera Infantum; decedents by sex and parentage, 37 years, (Table) Croup; statistics for 1892, and 37 years, (Table)	. 64
Croup; statistics for 1892, and 37 years, (Table)	65-67
Diarrings and Dysentery; recapitulation for 55 years	67-68
Diarrhea and Dysentery; recapitulation for 53 years.  Diphtheria; statistics for 1892, and 35 years, (Table).  Comparison of Diphtheria, Scarlet Fever and Croup.	. 71-72
Fever Malarial	72
Fever, Typhoid, 1892, and 37 years, (Table.)	74-78
Heart, diseases of; decedents by sex and parentage. 37 years, (Table.) Measles; statistics for 1892, and 37 years	. 79
Measles; statistics for 1892, and 37 years	83-84
Measies; Statistics for 1892, and 3; years.  Meningitis, Cerebro-spinal; deaths at each age, 37 years.  Phthisis; sex and parentage.  Phthisis; change in the rate of mortality.  Phthisis in Providence, 37 years, (Table).  Pneumonia; statistics for 1892, and 37 years, (Table).  Pneumonia and Bronchitis.	. 84-85
Phthisis; sex and parentage	- 88
Phthisis; change in the rate of mortality	. 91
Phthisis in Providence, 37 years, (Table)	. 89-90
Phetimonia; statistics for 1992, and of years, (Table.)	91-94
Scorlet Fever: statistics for 1802 and 38 years (Table)	95
Small-pox: recapitulatory table, 37 years,	99-91
Unknown causes	102
Still-born children in 1892	103-104
Scarlet Fever; statistics for 1892, and 38 years, (Table).  Small-pox; recapitulatory table, 37 years.  Unknown causes.  Still-born children in 1892.  Still-born children, 1856-1892.	. 105
GENERAL SUMMARY;	
Births, Marriages and Deaths; numbers and proportions, 38 years	100 100
Births, Marriages and Deaths; numbers and proportions, 38 years (Table XXI.).  Births and Deaths by parentage; excess of births, 37 years, (Table XXII.)	100-108
	ULL WLL



#### CITY REGISTRAR'S REPORT.

1892.

#### BIRTHS.

To the Honorable the City Council:

The great delay in preparing the Registration Report is due entirely to the difficulty in obtaining the record of births. The method by sending enumerators from house to house twice a year in accordance with the State law, has been shown to be defective, and this is supplemented by reports which are obtained from physicians attending births and from clergymen who baptise infants. This year these reports were not all obtained, compared, verified and arranged for recording until the first of May, so that the actual work of recording and making the tables could not be begun until that time. The enumerators are paid fifteen cents for each return and the clergymen and physicians are paid ten cents. In 1892 there were 3,527 returns obtained by the enumerators and 3,748 returns from physicians and clergymen. Many of these latter returns were duplicated in the different lists, but of these 3,748 births returned 479 were in addition to those returned by the enumerators. The total number of births occurring in the city of Providence during the year 1892, of which record has been obtained, is therefore 4,006, of which number 2,004 were males and 2,002 were females. This includes living births and still births. The total number gives a rate of 29.02 per thousand in a population of 138,000, or one in 34.47. There were also reported by the enumerators 169 births which occurred outside Providence. The parents of these children had recently become residents of the city and the statute requires that such births shall be recorded separately from those occurring in the city. These do not enter into the calculations of this report. In previous years such births have been not only recorded but reckoned in the tables, because it was believed that they no more than counterbalanced the number of children born in the city but

removed before the enumeration. The substraction of these 169 births makes the increase of the births in the city during 1892 over the number in 1891 seem less than it really is. Of the total number of 4,006 returns 62 were so imperfect or were obtained so late that they are not included in the tables. The total number entering into the tables is therefore 3,944.

In table I. is shown the number of children born in each ward, and in the whole city, in each month and quarter of the year. Also the number of children of each sex, in the city, and the number and sex of the colored children.

Table I. Births, 1892. Wards, Months, Sex and Color.

nwlin man	WARDS.									W	Colored children included in the preceding.					
Months.	I.	II.	III.	IV.	v.	VI.	VII	VIII.	IX.	X.	Males.	Females.	Total.	Males.	Females.	Total.
January February March	31 32 26	27 23 24	57 56 53	11 11 11	25 29 30	36 30 25	27 23 23	34 25 25	30 35 41	43 51 67	156 152 168	165 163 157	321 315 325	6 4 5	5 3 1	11 7 6
1st Quarter	89	74	166	33	84	91	73	84	106	161	476	485	961	15	9	24
April May June	34 30 28	28 32 39	53 67 50	12 7 12	23 25 23	35 21 30	26 22 20	20 27 27	28 37 42	49 59 47	159 155 169	149 172 149	308 327 318	6 6 8	4 7 2	10 13 10
2d Quarter	92	99	170	31	71	86	68	74	107	155	483	470	953	20	13	33
July August September	37 28 42	42 31 27	54 59 49	10 11 8	31 33 22	33 19 28	18 28 24	28 27 28	35 40 51	59 60 55	174 172 159	173 164 175	347 336 334	4 4 6	2 5 4	6 9 10
3d Quarter	107	100	162	29	86	80	70	83	126	174	505	512	1,017	14	11	25
October November December.	36 44 42	29 40 20	52 55 58	8 9 12	20 22 25	32 29 37	28 25 26	23 28 34	32 29 32	64 56 66	160 163 185	164 174 167	324 337 352	8 5 4	7 5 7	15 10 11
4thQuarter	122	89	165	29	67	.98	79	85	93	186	508	505	1,013	17	19	36
Wholeyear	410	362	663	122	308	355	290	326	432	676	1,972	1,972	3,944	66	52	118

The total number of births tabulated in 1892 was 7 less than in 1891; 92 less males and 85 more females.

In four wards of the city, the second, third, eighth and tenth, there were more births in 1892 than in 1891; in all the other wards the number was less than in the preceding year.

Ratio. The population of the city in 1892 was estimated at 138,000; the ratio of the 3,944 births, therefore, was 28.58 to each thousand inhabitants, or one in 34.99. For explanation of apparent decrease in rate see first page of report.

Season. The largest number of births in any one month in 1892 was 352 in December; the smallest was 308 in April; the largest quarterly total was 1,017 in the third quarter; the smallest 953 in the second quarter.

The following figures show the aggregate number of births in each quarter of the year for the thirty-six years, 1856 to 1891; the number in each quarter of 1892; and the same for the thirty-seven years, 1856 to 1892:

	Jan.	April.	First.	July.		Second	Whole
	March.	June.	6 mos.	Sept.	Dec.	6 mos.	Year.
Thirty-six years	19,940	. 20,684	40,624	.20,984 .	21,433.	42,417	83,041
1892	961	. 953	1,914	. 1,017	1,013	2,030	3,944
						-	
Thirty-seven years	20,901	.21,637	42,538	.22,001	22,446.	44,447	. 86,985

Proportion of the Sexes. The number of male children born in Providence in 1892 was 1,972; of female children, 1,972. The proportions of male and female children in each quarter and in the whole year were as follows:

	Males.	Female	s. Male	s. Fems	les.Males I	emales.	
JanMarch	476	485, 0	or 98.1	to 100,	or 49,53 and	50.47 in e	ach 100 born.
April-June	483	470, 0	or 102.8	to 100,	or 50.68 and	49.32 in ea	ach 100 born.
July-Sept	505	512,	or 98.6	to 100,	or 49.66 and	50.34 in e	ach 100 born.
OctDec	508	505, 0	or 100.6	to 100,	or 50.15 and	49.85 in ea	ach 100 born.
				_			
Whole year	,972	.1,972, 0	or 100.0	to 100,	or 50.00 and	50.00 in e	ach 100 born.

The following table shows the same facts for the thirty-seven years, 1856–1892, inclusive:

The following table shows the relative proportion of the sexes for the thirty-nine years, 1854 to 1892; it is given by periods of five years, from 1854 to 1888. The whole number of children born during the thirty-nine years was 89,954:

Table II. Births, 1854 to 1892. Proportion of the Sexes.

	Males.	Females	. Males.	Females.	Males. F	'emales.		
1854-1858	4,171	3,885, 01	107.4 to	100; or	51.77 and	48.23 in	each 100	born.
1859-1863	4,050	3,752, 01	: 107.9 to	100; or 8	51.91 and	48.09 in	each 100	born.
1864-1868	3,977	3,787, 01	: 105.0 to	100: or	51.22 and	48.78 in	each 100	born.
1869-1873	5,056 .	4,946, or	102.2 to	100; or 5	50 55 and	49.45 in	each 100	born.
1874-1878	6,817	6,428, 01	106.0 to	100; or 8	51.47 and	48.53 in	each 100	born.
1879-1883	7,037	6,619, 01	106.3 to	100: or 8	51.53 and	49.47 in	each 100	born.
1884-1888	7,681	7,235, 01	106.1 to	100; or 8	51.49 and	48.51 in	each 100	born.
1889	1,592 .	1,579, 01	100.8 to	100; or 8	50.20 and	49.80 in	each 100	born.
1890	1,698	1,719, or	98.8 to	100; or	49.69 and	50.31 in	each 100	born.
1891	2,064.	1,887, 01	r 109.3 to	100; or	52.24 and	47.76 in	each 100	born.
1892	1,972	.1,972, or	100.0 to	100; or 8	50.00 and	50.00 in	each 100	born.
39 years	46,115	43,839, or	105.2 to	100; or 5	51.27 and	48.73 in	each 100	born.

The statistics for each year from 1854 to 1882, inclusive, will be found in the report for 1882, page 5; for 1883, in the report for that year, page 3; for the years 1884 to 1888, inclusive, in the report for 1889, page 3.

Color. The number of colored children born in 1892 was reported as 118; this was 15 more than in the preceding year. Considering the colored population to bear the same proportion to the whole population of the city as in 1890 we find it to be 4,237; of births in this class, the ratio was one in 35.90, or 27.85 births to each 1,000 colored inhabitants.

The colored children were born in different wards, as follows: First, 17; second, 23; third, 5; fourth, 6; fifth, 9; sixth, 9; seventh, 21; eighth, 24; ninth, 4; tenth, 0.

The number in each month and quarter was:

January11 February7 March6	April 10 May13 June10	July 6 August 9 September10	October15 November10 December11
1st Quarter24	2d Quarter 33	3d Quarter25	4th Quarter36

There were 66 males and 52 females among the colored children born in 1892, a proportion of 126.9 males to each 100 females; or 55.93 males and 44.07 females in each 100 children.

The births, marriages and deaths among the colored population are given in the next table, in periods of five years, from 1856 to 1890, for 1891 and 1892, and for the whole period of thirty-seven years. The figures for the single years may be found in the reports for 1882, 1885 and 1890:

	Births.	Marriages.	Deaths.
1856-1860	222		267
1861-1865	228	117	291
1866-1870	421		365
1871-1875	512	257	543
1876-1880	511	238	529
1881-1885	565	273	558
1886-1890	581	302	632
1891	103	57	130
1892	118		143
		66 contractions	
37 years	3,261		3,458

The proportion of deaths to births among the colored population in 1892 was 121.2 to 100; in the thirty-seven years, 1856 to 1892, the proportion was 106.0 deaths to each 100 births.

#### PARENTAGE.

The following tables give statistics as to the parentage of the children born in the city; the facts are interesting and important, as showing the variety of nationalties represented and the proportion of each class.

Table III. shows how many children of each class were born in each ward of the city, and the number and percentage of each class in the whole city. The parentage is reckoned according to the birthplace of the father.

Table III. Births, 1892. Wards and Parentage.

BIRTH-PLACES OF				7	W A	RD	s.				Whole City.	In each 100 born there were—
FATHERS.	I.	II.	111.	IV.	v.	VI.	VII.	vIII.	IX.	х.	Whol	In ea borr
America (mother Amer'n)	99	146	135	50	107	127	146	124	108	153	1,195	30.30
America (mother foreign)	41	35	60	14	38	54	22	36	35	70	405	10.27
Foreign countries (mother American)	33	36	48	16	29	29	24	32	35	83	365	9.25
Ireland	91	46	132	23	72	45	24	39	70	132	674	17.09
Great Britian	8	20	54	1	7	56	26	20	28	128	348	8.82
Germany	9	2	17	2	13	3	15	7	3	6	77	1.95
British America	4	11	8	2	5	1	8	4	4	12	59	1.50
French Canada		4	12	2	6	3	7	40	5	53	132	3.35
Portugal & Western Islands	111						1	1	1		114	2.89
Sweden	2	3	9	1	11	26	3	4	4	18	81	2.05
Norway	1	1	1	2	2	4	2	1		2	16	-40
Italy	1	1	52	4	2		2	3	129	10	204	5.17
Russia	2	30	65		6	2	2	2	1	2	112	2.84
Poland	1	19	4				1	2		2	29	-73
Austria		3	4		2			1			10	-25
Hungary		2			1			1			4	-10
Roumania					]	]					2	-05
France			2		1		2		2		7	-18
Spain	1										1	.03
Denmark						1				2	3	.07
Switzerland			1		2		1	1			5	.13
Syria									2		2	.05
Arabia									1		1	•03
Asia·····									1		1	-03
Bahama Islands	1										1	•03
Atlantic Ocean									1	1	2	-05
Total Foreign	232	142	361	37	131	142	94	126	252	368	1,885	47.79
America (mother unkn'n)				1							1	.03
Foreign (mother unkn'n)				1							1	-03
Unknown (mother Am'n)	2	2	29	2	1	1	4	7	2	2	52	1.32
Unknown (mother foreign)	2		29		1	2		1			35	-88
Both unknown	1	1	1	1	1						5	-13
Ward totals	410	362	663	122	308.	355	290	326	432	676	3,944	100.00

The statement given below shows the number and percentage of children of American, foreign and mixed parentage since 1856. From 1856 to 1890 the statistics are given by periods of five years; then they are given for 1891 and 1892, and for the whole period of thirty-seven years.

The figures for 1891 and 1892, however, do not include the whole number of births for those years, as there were, in 1891, 82 cases, or 2.08 per cent. of all, and in 1892, 94 cases, or 2.39 per cent., where the birthplace of one or both parents was not stated. The same is true of the total for thirty-seven years; the 176 cases being 0.20 per cent. of the whole number of births during that period:

			Am. father.	For, father.
	American.	Foreign.	For, mother.	Am. mother.
Years. Nu	imber. Perci. N	umber. Per ct.	Number. Per ct.	Number. Per ct.
1856-1860	3,32739.95	4,42453.10	292 3.51	285 3.42
1861-1865	2,79637.57	3,80952.88	352 4.89	336 4.66
1866-1870	3,70241.92	4,11846.64	447, 5.06	563 6.38
1871-1875	4,722 40.23	5,35345.61	693 5.90	970 8.26
1876-1880	5,17339.95	5,59743,22	929 7.17	1,251 9.66
1881-1885	5,87640.81	5,78640.18	1,215 8.44	1,52210.57
1886-1890	5,88437.62	6,68642.74	. 1,474 9.42	1,59810,22
1891	1,32733.59	1,79645.45	365 9.24	381 9.64
1892	1,19530.30	1,88547.79	40510.27	365 9.25
`—				
37 years3	3,91238.99	39,45445.36	6,172 7.10	7,271 8.36

The percentage of children of American parentage was less in 1892 than in any preceding year, and the percentage of children of American parentage for the five years, 1886 to 1890, was less than for any period of five years, except the war period from 1861 to 1865.

For fuller particulars for each year, see reports for 1882, page 9; 1885, page 7, and 1890, page 8.

The percentage of children born in Providence of parents of each nation is as follows, for each of the thirty-seven years, 1856-1892, inclusive:

		English,	British	French	ı S	wedish &	\$z	Other
American.			man. Am.	Can.	Portu.	Norw'n.	Italian	For'n
		Per ct. Per	ct. Per ci	t. Per ct.	Per ct.	Per ct.	Per ct.	Per ct.
185642.09	41.79	4.842.	27					2.27
185738.51	44.32	. 5.923.	73					1.54
1858 41.30	42.98 .	4,822.	26					1.33
185937.60	45.58	5.343.	26					1.51
186040.11	40.96	6.552.	85					1.64
186137.91	42.61	6.032.	49					1.97
186238.52	41.53	6.02 3.	07					1.64
1863 35.73	43.08	6.35 2.9	98					1.99
186438.69	41.36	. 6.552.	75					1.34
186536.67								1.93
186640.20			21					2.32
186739.72	37.81 .	5.663.0	)2					2.21
186841.75	36.01	6.481.8	881.07					1.07
186945.19	34.14	4.01 1.8	311.32					1.38
1870 42.35			961.91.					90
187144.85			332.09					1.73
187241.12								2.31
187340.84								2.16
187438.42					1.22			1.19
1875 37.50					98			1.43
187638.27					1.22			1.25
187741.45					1.10			1.30
		. 6.742.8			. 1.18			1.71
187940.80					96			1.70
188039.90					1.60			1.33
188140.71					1.07			2.43
188241.10					1.08			2.98
		9.022.6			79			2.23
188441.00					1.25			3.35
188540.56			53.54		1.64			3.27
188640.71					1.66			4.49
188737.56						1.04		
		9.201.7				1.53		
		9.78 2.3				1.52		
		. 10.151.7				1.99		
1891 33.59	16.17 .	9.191.9	01.57	3.67 .	2.17	2.79	4.50	3.49
189230.30	17.09 .	8 82 1.9	51.50	3.35	. 2.89	2.45	5.17.	4.57

The British-American were included in the "other foreign" until 1868; the Portuguese until 1874; the Swedish, Italian and Norwegian until 1887. In 1891 the French-Canadians were first reckoned separately from natives of the other British Provinces.

Table IV. shows the parentage of the children more exactly by giving the birthplace of both parents:

Table IV. Births, 1892. Parentage.

		BIF	RTH-	-PI	AC	ES (	)F N	10	TI	IER	s.		ord-
BIRTH-PLACES OF FATHERS.	United States.	Ireland.	Great Britain.	Germany.	British America.	French Canada.	Portugal and Western Islands.	Sweden.	Norway.	Italy.	Other Countries.	Unknown.	Whole number according to the father's birth-place.
United States	1,195	161	114	9	80	28	1	4	1		7	1	1,601
Ireland	131	610	45		16	1		1			1		805
Great Britain	112	52	280		14			1			1		460
Germany	24	8		66	2			!			1		101
British America	32	13	4	2	39						1		91
French Canada	33	4			1	127							165
Portugal and Western Islands	5	1					113		٠.				119
Sweden	1	2				1		76	1		1		82
Norway	2	1			1	1		1	12				18
Italy	8	1	1							201	1	1	213
Other Countries	17	3	4	2		1	1				169		197
Unknown	52	15	5	4	5	2	1	3				5	92
				-	_							-	
Whole number according to the mother's birthplace	1,612	871	453	83	158	161	116	86	14	201	182	7	3,944

The 1,195 children whose parents were both born in the United States were 30.30 per cent. of all the children born during the year; this was 132 less children of this class than in 1891, and 3.29 per cent. less. There were, in addition to these, 405 children whose fathers were born in the United States and the mothers in some foreign country; and 1 whose father was born in this country, the mother's birthplace being unknown; there were, therefore, 1,601 children of American fathers, or 40.59 per cent. of all. There were also 365 children whose fathers were foreign and the mothers American; and 52 where the mother was American, the father's birthplace being unknown; a total of 1,612 children of American mothers, or 40.87 per cent. of all. The

children of the two classes of mixed parentage numbered 770, or 19.52 per cent. of all the children born during the year; 24 more children and 0.64 per cent. more than in 1891. The children of purely foreign parentage in 1892 numbered 1,885, or 47.79 per cent. of all; 89 more children, and 2.34 per cent. more than the same class in 1891.

#### NUMBER OF THE CHILD.

Table V. shows how many of the children born in 1892 were the first, second, third, etc., children of their mothers; also the same statistics for the 88,585 children born during the thirty-eight years, 1855-1892:

NU	MBEF	2 0	F CH	ILD.		1892.		38 Years
					American.	Foreign.	Total.	1855-1892
First	child	of	moth	er	519	558	1,077	22,190
Second	6.6	6.6	4.6		326	487	. 813	17,920
Third	6.6	. 6	4.6		239	376	615	13,875
Fourth	4.5	6.6	4.6		166	282	448	10,432
Fifth	< 6	4.6	6.6		119	202	321	7,706
Sixth	6.6	4.6	4.6	*(* * * * * * * * * * * *	91	113	. 204	5,581
Seventh	6.6	6.6	4.6		55	103	158	3,964
Eighth	4.5	6.6	6.6	*********	46	70	116	2,674
Ninth	6.6	6.6	6.6		25	52	77	1,728
Tenth	4.6	6.6	4.6		12	27	39	1,152
Eleventh	4.6	6.6	6.6		5	32	37	628
Twelfth	6.6	6.6	4.6		5	11	16	382
Thirteenth	6.6	4.6	6.6		2	4	6	165
Fourteenth		6.6	4.6		1	1	2	85
Fifteenth	6.6	6.6				3	3	41
Sixteenth	6.6	6.6	6.6	* * * * * *	1	1	. 2	16
Seventeenth	6.	6	6.6			1	1	14
Eighteenth	6.6	66	6.6	* * * * * * * * * * * * * * * * * * * *		1	1	5
Nineteenth	6.6	6.6	6.6					4
Twentieth	٠.	6.6	66					1
Twenty-third	6.6	6.6	6.6			1	1	2
Statistics im	oerf <b>e</b> c	t, n	ot tab	ulated	•••••		7	20
Whole numb	er of	chil	dren.		1,612	2,325	3,944	88,585

In 1891, for the first time, the statistics as to the number of the child and the age of the mother were distinguished according to parentage, but there were nine cases where the nativity of the mother was not stated; also there were four cases, two of Amercan mothers and two of foreign mothers, where the number of the child was not stated. In 1892, the nativity of seven mothers was unknown. In all the tables on this subject, therefore, the statistics are imperfect, and the number of children does not correspond with the total number born.

The number of *first* children in 1892 was 1,077, or 27.28 per cent. of the 3,944 children born during the year; this was 40 more children and 1.03 per cent. more than in 1891. In the thirty-eight years, 1855 to 1892, there were 88,585 children born; of this number, 22,190, or 25.05 per cent. were the first children of their mothers.

Average Number of Children to Each Mother. There were forty-two twin births in 1892; the number of mothers, therefore, was 3,902, but in tables VI. and VII. the number of mothers for 1892 is given as only 3,895 (because of the defective statistics mentioned above); these 3,895 mothers had borne in all 12,979 children, an average of 3.33 children to each mother; the average number of children to American mothers was 3.07; to foreign mothers, 3.51.

The following table shows the average number of children to each mother in the thirty-eight years, 1855-92; the statistics are given by periods of five years from 1855 to 1889; for 1890, 1891 and 1892 they are given by parentage so far as this could be determined:

Table VI. Births, 1855 to 1892.

Average Number of Children to Each Mother.

Vonna		No. of			
rears.	Motners.	Children. No.	of Children. Aver	age.	
1855-1859	8,186	8,280	. 27,8153.40	children to	each mother.
1860-1864	7,456	7,553	. 26,784 3.59	children to	each mother.
1865-1869	8,126	8,239	. 28,158 3.47	children to	each mother.
1870-1874	10,812	10,963	. 35,6023.29	children to	each mother.
1875-1879	12,850	12,987	. 44,955 3.50	children to	each mother.
1880-1884	13,937	14,090	49,8813.58	children to	each mother.
			. 52.356 3.49		
1890 J Am	1,496	1,522	4,741 3.17 6,716 3.60	children to	each mother.
For	1,864	1,895	. 6,7163.60	children to	each mother.
1901 J Am	1,741	1,748	5,359 3.08 . 7,650 3.52	children to	each mother.
1892 [ Am	1,600	1,612	. 4,9133.07 . 8,0663.51	children to	each mother.
/ For	2,295	2,325	. 8.0663.51	children to	each mother.
26 700 20	27 590	99 560	302,9963.46	abildman to	oooh mothon
oo jeals	01,020	** 00,000	00%,000	CHIMOLEH 10	rach mother.

Table VII. Births, 1892. Number of Child with Age of Mother.

American Mothers.

The state of the	Mothers			NI	JMB	ER	OF '	THI	E C	H	IL	D.					rs at		e in	they had	of rs at
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	OF THE YEARS.	First.	Second.	Third.	Fourth.	Fifth.	Sixth.	Seventh.	Eighth.	Ninth.	Tenth.	Eleventh.	Twelfth.	Thirteenth.	Fourteenth.	Sixteenth.	Number of Mothers at	each Age.	Children they bore 1892.	Total Children they borne.	n ase
Totals   519   326   239   166   119   91   55   46   25   12   5   5   2   1   1   1,600   1,612   4,913   3.07	15.	1 8 8 8 19 *35 8 45 51 52 50 *41 1 26 *33 15 14 1 12 2 4 4 6 6 2 2 2 3 3 3	10 8 15 266 288 333 411 *299 233 233 112 233 24 11	111 188 144 166 17 189 181 181 181 181 181 181 181 181 181	10 6 122 14 11 11 11 14 14 14 15 6 6 2 2 2 1 2	4 4 4 10 12 12 12 12 17 7 13 8 8 8 5 9 9 4 3 3 · · · · · · · · · · · · · · · · ·	1 *1 *4 *4 *5 5 3 10 12 12 9 8 8 6 6 10 6 5 4 4 1 3 3 1 · · · · · · · · · · · · · · · ·	3 2 1 2 5 4 4 4 8 *10 2 2 3 4 4 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	· · · · · · · · · · · · · · · · · · ·	\$\frac{4}{5}\$\frac{1}{1}\$\frac{1}{2}\$\cdots\$\frac{1}{1}\$\cdots\$\c	3 1 2 1	1	1 1	1	1		3	1 8 10 22 457 68 91 08 04 14 04 86 86 57 87 57 63 57 63 57 63 13 11 41 41 41 41 41 41 41 41 41 41 41 41	1 1 8 10 222 466 477 688 104 417 105 588 577 633 599 366 25 199 133 111 4 1	1 1 8 8 8 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9	1.00 1.00 1.20 1.14 1.27 1.32 1.48 1.64 1.87 1.81 2.16 2.13 3.42 3.63 3.12 3.63 3.81 4.32 4.65 5.00 5.08 5.91 5.67 4.58 6.00 5.08 13.00 8.00 1.00

<sup>\*</sup> One of these mothers bore twins.

Table VII. Births, 1892. Number of Child with Age of Mother. FOREIGN MOTHERS.

16. 2 17. 7 8 18. 17 4 19. 221 11. 20. 38 24 21. *38 21. 22. *58 27 23. 55 29 24. *443 39. 25. 46 *42. 26. *33 44. 27. 29 *44 *27. 29 *44 *28. *51 *47 *28. *51 *47 *47 *28. *51 *47 *47 *28. *51 *47 *47 *48 *51 *47 *48 *51 *48 *51 *47 *48 *51 *51 *48 *51 *51 *51 *51 *51 *51 *51 *51 *51 *51	Second. Third.	0 0								Mothers	bore	they	Chil
18. 17 4. 17 1. 19. 21 11. 20. 38 24 21. 21 12. 22. *58 27 23. 55 29. 25. 46 *42 26. *63 44 27. 29 *44 28. *51 *47 29. 19 25 30. 29 38 31. 4 *17 32. 12 18 33. 5 14 34. 1 13 35. 6 †9 36. †4 8 37. 4 4 8 38. 2 39. 1 3 39. 1 3	Sec	Fifth.	Seventh.	Ninth.	Tenth. Eleventh.	Twelfth.	Fourteenth.	Sixteenth.	Eighteenth. Twenty-third.	Number of Meach Age.	Children they 1892.	Total Children they had borne.	Average No. of Children to Mothers at each Age.
41 1 42 43 44 45 46 47 53 Not given 2 1	4	31 *15 (40 17	4 6 3 1 6 2 6 13 5 7 *5 4 9 3 2 10 5	2 2 1 2 4 1 7 3 8 6 3 8 6 1 1 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1	1 1 2 2 2 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4				1	2 111 211 33 66 65 104 106 117 151 155 166 198 174 173 174 173 174 175 175 176 177 177 177 177 177 177 177 177 177	2 111 211 33 66 66 66 105 116 118 152 157 77 170 170 175 170 175 174 112 193 78 74 44 44 44 44 44 45 46 46 46 46 46 46 46 46 46 46 46 46 46	30 19 18	1.00 1.45 1.15 1.39 1.68 1.54 1.68 1.59 2.18 3.02 2.18 3.02 2.18 4.13 4.13 4.13 4.13 4.13 4.13 4.13 4.13
Totals 558 487 8			1							1 4	1	8	1.75

The reference marks in this table indicate the number of the first child in cases of twin births; thus of American mothers at the

<sup>\*</sup>One of these mothers bore twins. †Two of these mothers bore twins.

age of 19, one mother bore twins which were her first and second children; of foreign mothers at the age of 27, four mothers bore twins, in one case the second and third children of the mother, in two cases the third and fourth children, and in one, the fifth and sixth children.

The number of mothers in 1892 who were under 20 years of age was 154; 87 American and 67 foreign, or 3.96 per cent. of mothers of known age. This was 0.57 per cent. more than the average for twenty-eight years and six months, the time during which a record of the ages of the mothers has been kept.

The number of mothers in 1892 who were 40 years old and over was 175; 51 American and 124 foreign, or 4.50 per cent. of mothers of known age. This was 0.48 per cent. less than the average for twenty-eight years and six months.

The following table shows the number and percentage, at different age-periods, of the 73,851 mothers whose ages have been recorded since July 1, 1863:

	Number of mothers.	Per cent
Under twenty years	2,507	3.3
Twenty, and under twenty-five		21.8
Twenty-five, and under thirty		
Thirty, and under thirty-five		
Thirty-five, and under forty		
Forty, and under forty-five		
Forty-five and over		
		40-00-00-
Total		100.0

The average age of these 73,851 mothers is 28.92 years. This is 0.31 years more than the average age of the mothers of 1892, which was 28.61. The average age of American mothers in 1892 was 28.07 years; of foreign mothers 28.99 years. The average age of the mothers for the period above mentioned is 5.67 years more than that of the brides for the last 37 years. The average age of the mothers for 1892 is 3.15 years more than that of the brides of the same year.

During this period of twenty-eight years and six months, the number of children borne by mothers whose age was stated, was

74,613, the number of children to mothers at each age was as follows:

ge of mother. Children.	Age of mother. Children	. Age of mother. Children
13	274,561	41 55
14 7	28 5,214	42 63
15 27	293,935	43 36
16 105	305,685	44
17 319	312,659	45
18 758	323.637	46
191,298	<b>3</b> 33,023	47
20 2,043	342,618	48
212,370	353,518	49
223,481	36 2,305	50
233,983	371,905	51
244,378	382,072	53
255,001	39	54
264,629	40	
All known ages		

The 74,613 children were divided as follows, to mothers of different age periods:

	Number of mothers.	Per cent
Under twenty years		3.37
Twenty, and under twenty-five		21.79
Twenty-five, and under thirty		
Thirty, and under thirty-five		23.62
Thirty-five, and under forty		
Forty, and under forty-five		
Forty-five and over		
Total		100.00

#### PLURALITY BIRTHS.

The number of twin births in 1892 was 42; 12 more than in 1891. Cases, 42; children, 84.

Locality. First ward, 6; second, 6; third, 8; fourth, 2; fifth, 3; sixth, 3; seventh, 1; eighth, 2; ninth, 3; tenth, 8.

Season.

January 8 February 1 March 3	April	July 3 August 2 September 5	October 7 November 3 December 2
1st quarter12	2d quarter8	3d quarter10	4th quarter12

Sex. In 18 cases both children were males; in 6 both were females; and in 18 there was one child of each sex. In all there were 54 male, and 30 female children.

Color. In 1 case the parents were colored; all others white.

Parentage. The parentage of the 42 cases of twin births was as follows:

Cases.	Parentage.	Cases.	Parentage.
9	American.	1Po	olish.
6	Irish.	3A	merican and Irish.
1	Scotch.	3A1	merican and English.
4	Portuguese.	2Ir	ish and Amercian.
2	French-Canadian.	1 A	merican and British American
1	Swedish.	1 Go	erman and American.
1	Norwegian.	1Ir:	ish and British American.
	Italian.	1 Sv	viss and Scotch.
1	Russian.		

#### THIRTY-EIGHT YEARS' SUMMARY OF PLURALITY BIRTHS.

In the thirty-eight years, 1855 to 1892, there were 1,024 cases of plurality births in Providence; 1,014 of twins; 9 of triplets; and 1 of quadruplets.

Sex. Of the nine cases of triplets, in one case, all the children were males; in five cases, all were females; and in three cases there were two male children and one female; making 9 male and 18 female children in the nine cases.

In the case of the quadruplets, all the children were males.

In the 1,014 cases of twins, there were 341 where both children were males; in 341 both were females; and in 332 cases there was one male and one female child; in all 1,014 male, and 1,014 female children.

In the 1,024 cases of plurality births, there were 1,027 male and 1,032 female children; a proportion of 100.5 females to 100 males; or 50.12 female and 49.88 male children in each 100 born.

Color. In the cases of triplets and quadruplets, the parents were all white; of the 1,014 cases of twins, in 946 the parents were white, and in 68 they were colored.

Parentage. Of the triplets, in one case the parents were both American; in five cases, both were Irish; in one case, both Ger-

man; in one the father was English and the mother Irish; and in one, the father Scotch and the mother Irish. There was but one American mother in the nine cases.

Of the quadruplets, the father was American, and the mother British-American.

The parentage of the 1,014 cases of twins is shown in the following table, the first letter indicating the birth-place of the father, the second that of the mother:

A., American; I., Irish; E., English; S., Scotch; W., Welsh; G., German; B. A., British-American; F. C. French-Canadian; D., Dutch; P., Portuguese; It., Italian; Sw., Swedish; N., Norwegian; F., French; Br., Brazilian; R., Russian; Pol., Polish; W. I., West Indian; Aus., Austrian.

Parentage. No.	Parentage. No.	Parentage. No.	Parentage. No.
A. and A330	Pol. and Pol 3	Sw. and I 1	E. and S 2
I. and I307	Br. and Br 1	It. and I 2	E. and B. A 4
E. and E 45	I. and A 35	P. and I 1	I. and W 2
S. and S 17	E. and A16	A. and E11	A. and B. A 10
G. and G 28	S. and A 3	I. and E14	I. and B. A 4
B. A. and B. A. 25	B. A. and A10	S. and E 1	F. and B. A 1
F. C. and F. C 2	G. and A 3	G. and E 1	S. and B. A 1
D. and D 1	Sw. and A 2	B. A. and E 1	A. and Sw 1
N. and N 3	A. and I41	A. and S 5	Aus. and R 1
P. and P 22	E. and I15	F. and A 1	Swiss and S 1
It. and It 13	S. and I 3	W. I. and A 1	
Sw. and Sw 3	I. and G 2	N. and E 1	Total,1,014
P. and A 1	G. and I 1	It. and F 1	
R. and R 4	B. A. and I 4	I. and S 6	

In the 1,014 cases of twin births, 402 mothers, or 39.64 per cent. were American; 375, or 36.98 per cent., were Irish; and 237, or 23.37 per cent. were of other foreign nations. In the 1,024 cases of plurality births, 403 mothers were American; this was 0.97 per cent. of the 41,570 American mothers who bore children during the 38 years, 1855 to 1892; 621 mothers were foreign, which was

1.36 per cent. of the 45,664 foreign mothers who bore children during the same period.

Proportion of plurality births to whole number. The following figures show the number and proportion of plurality births in 1892 and in the thirty-eight years, 1855-1892:

	Single Births.	Twin.	Triplet. Quae	Whole No. druplet. of Births.	Whole No. of Children.
				183.648 . 0 3,902 .	
38 years, 1855-	1892 86,526	1,014	9	1	88,585

#### Proportion of cases of plurality births:

185 <b>5-</b> 1891		1 in 9,294.2	·
1855-1892	1 in 86.3	1 in 9,727.7	1 in 87,550

#### Proportion of children of plurality births:

1855-1891	in 43.51	Triplets. in 3,134.8	·
-			
1855 18921	in 43.7	in 3,280.9	1 in 22.146.2

Proportion of plurality births by parentage. In the thirty-eight years, 1855 to 1892, the whole number of births was 87,550. The following table shows the number of births of American, foreign and mixed parentage, and the number and proportion of plurality births in each of these classes:

Whole	No. Births.	Plurality Cases.	1 in	Per ct.
American parents	. 34,219	331	103.6	0.97
Foreign parents	.39,749	552	. 72.0	1.39
American father, foreign mother.	6,147	69	89.0	1.13
Foreign father, American mother.	. 7.259	72	.100.8	99
Total	.87,374	1,024	. 85.3	1.17

Of the 1,024 cases of plurality births during the thirty-eight years, the 331 of American parentage were 32.32 per cent.; the 552 of foreign parentage were 53.91 per cent.; the 69 with American fathers and foreign mothers, 6.74 per cent.; and the 72 with foreign fathers and American mothers, 7.03 per cent.

Illegitimacy. The number of illegimate births recorded in Providence in 1892 was 65, or 1.65 per cent. of all the children born during the year. The actual number was undoubtedly greater than this, as the method of obtaining records by enumerators is less likely to secure accurate information on this than on any other point.

Thirty-six of these births occurred at the Providence Lying-In Hospital.

In 54 cases this was reported as the first child of the mother, in five cases the second child, in two cases the third, in one case the fifth, and in two cases the sixth, and in one case the eighth child.

#### The nativity of the mothers was as follows:

United States40	Sweden	
Ireland11	England	1
Canada and British Provinces 5	Germany	2
Western Islands 1	Not stated	2

The 40 mothers of American birth were 61.54 per cent. of all. Of these 22, or 33.85 per cent. of all, were born in Providence. Thirteen of the mothers, or 20.00 per cent. were colored.

#### The ages of the mothers were as follows:

No. of		No. of		No. of
Age. mothers.	Age.	mothers.	Age.	mother
4 1	21		28	****
5 1	22	5	30	
6 3	23	5	34	
7 5	24	2	35	
87	25	4	36	
99	26	2	38	
04	27	1		-
Total			• • • • • • • • • • • • • • • • • • • •	6

#### MARRIAGES.

The number of marriages reported in Providence for 1892 was 1,592, which was 104 more than in 1891.

The following table shows the number of marriages in each month and quarter of 1892, and each of the six years preceding; also the aggregate and average numbers for each month and quarter of the thirty-eight years, 1855 to 1892:

TABLE VIII. MARRIAGES. Season.

MONTHS.	1892.	1891.	1890.	1889.	1888.	*1887	1886.		38 Years' Average.
January	127	120	115	119	124	133	99	3,284	87
February	127	91	106	101	108	108	82	2,878	76
March	82	63	59	81	39	64	91	1,877	49
lst quarter	336	274	280	301	271	305	272	8.039	212
April	126	173	155	118	132	118	67	3.158	83
May	109	92	105	105	110	104	99	3,093	81
June	195	175	150	140	138	27	125	3,599	95
2d quarter	430	440	410	363	380	349	291	9,850	259
July	96	99	93	90	94	80	92	2,695	71
August	118	114	105	<b>1</b> 06	106	76	99	2,546	67
September	149	152	127	105	117	126	128	3,401	89
3d quarter	363	365	325	301	317	282	319	8,642	227
October	183	163	165	147	149	144	143	3,956	104
November	169	155	152	160	162	159	135	4,180	110
December	111	91	98	95	70	97	89	2,849	75
4th quarter	463	409	415	402	381	400	367	10.985	289
Yearly totals	1,592	1,488	1.430	1,367	1,349	1,336	1,249	37,516	987

<sup>\*</sup>After the report for 1877 was published, 52 marriages were added to the records for that year; the figures have, therefore, been corrected in this table, and in all tables where the year 1887 is included.

The population of the city, in 1892, was estimated at 138,000; the number of persons married during the year was 3,184, a ratio of one in 43.34 of the population, or 23.72 in each thousand. The ratio of persons married to births, in 1992, was 87.30 to 100; of persons married to deaths, 107.4 to 100.

Season. As a rule the largest number of marriages in any one month is in November, but in 1892 it was in June, the number being 195. The smallest was 82 in March. The largest quarterly total was 463 in the fourth quarter.

The following figures show the number and per cent. of marriages in each quarter of 1892, and of the thirty-seven years, 1855 to 1891:

	189	2.	1855-1	1855-1891.				
Mε	arriages.	Per cent.	Marriages.	Per cent.				
JanMarch	336	21.11	7,703	21.44				
April-June	430	27.01	9,420	26.22				
July-Sept	363	22.80	8,279	23.05				
OctDec	463	29.08	10,522	29.29				
Year	1,592	100.00		100.00				

Table IX. shows the birthplaces of the 3,184 persons married in 1892:

Table IX. Marriages, 1892. Nativity of the Parties.

		BI	RTH	-PL	ACE	es o	F B	RID	ES.		ms.
BIRTH-PLACES OF GROOMS.	United States.	Ireland.	England, Scotland and Wates.	Germany.	British America.	French Canada.	Portugal and Western Islands.	Sweden and Norway.	Italy.	Other Countries.	Whole No. of Grooms
United States	745	58	49	2	32	5	2	4		3	900
Ireland	39	186	11		3						239
England, Scotland and Wales	50	9	75		4			3		3	144
Germany	10	4		17	1	1				4	37
British America	28	4	3		16						51
French Canada	15	2	1		1	21					40
Portugal and Western Islands	5	2					39				46
Sweden and Norway	5	2		2	1			46			56
Italy	7		1		1				38		47
Other Countries	9	2		2						19	32
							_				
Whole number of Brides	913	269	140	23	59	27	41	53	38	29	1,592

Bride and groom born in the same country. There were 745 marriages in 1892 where both bride and groom were natives of the United States; 46.80 per cent. of all the marriages during the year. This was 72 more marriages and 1.57 per cent. more than in 1891. In 186 marriages, or 11.68 per cent., both parties were natives of Ireland; 43 more marriages and 1.68 per cent. more than in the preceding year. In 75 marriages both parties were natives of Great Britain; this was 14 less than in 1891; in 17 both were born in Germany; in 21 cases both were French-Canadians, and in 16 both were natives of other British Provinces. In 39 cases both were born in Portugal or the Western Islands, in 46 both in Sweden or Norway, and in 38 both in Italy. There were 19 marriages in which both parties were natives of countries not named in the table; some, but not all of them, marrying natives of the same country.

Bride and groom born in different countries. The whole number of grooms born in the United States was 900; of these 155, or 17.22 per cent., married foreign women. Of the 913 brides born in the United States, 168, or 18.40 per cent., married men of foreign birth. Of the 239 grooms born in Ireland, 53, or 22.17 per cent., married women of some other country; of the 269 brides born in Ireland, 83, or 30.85 per cent., married natives of some other country than their own. There were 144 grooms who were natives of Great Britain; 69 of them married women of other nationalities.

There were 32 grooms born in countries not named in the table. Their nativity was as follows: Russia 11, Austria 4, Poland 4, St. Helena 2, Denmark 2 and Alsace, Asia, Guatemala, Holland, Hungary, Isle of Jersey, Roumania, Switzerland and Syria 1 each.

There were 29 brides born in countries not named in the table. Their nativity was as follows: Russia 8, Austria 5, Poland 5, Switzerland 3, Denmark 2, Africa, Brazil, France, Holland, St. Helena and Syria 1 each.

The following table shows the number and percentage of American, foreign and mixed marriages in the thirty-eight years, from 1855 to 1892; the statistics are given for periods of five years from 1855 to 1889; and for the years 1890, 1891 and 1892. The same statistics for each one of the twenty-eight years, 1855–1882, will be found on page 25 of the Report for 1882:

	Ame	rican.	For	eign.	Mixed.
Years.	Number.	Per cent.	Number.	Per cent.	Number. Per cent.
1855-1859	1,660	54.25	1,090	35.62	31010.13
1860-1864	1,697	55.40	989	32.29	37712.31
1865-1869	2,433	58.23	1,216	29.11	52912.66
1870-1874	2,966	55.84	1,50	28.39	83815.77
1875-1879	2,932	57.85	1,256	24.78	88017.37
1880-1884	3,254	54.90	1,449	24.45	1,22420.65
1885-1889	3,166	49.53	1.890	29.57	1,33620.90
1890	646	45.17	463	32.38	32122.45
1891	673	45.23	506	34.00	30920.77
1892	745	46.80	524	32.91	32320,29
				-	
38 years	20,172	53.78	10,891	29.03	6,44717.19

The whole number of marriages given in the preceding table for thirty-eight years is 37,510, while in Table VIII. it is 37,516; this is owing to imperfect statistics in the early records. See Report for 1879, page 22.

Marriages of colored persons. There were 60 marriages in 1892 in which both parties were colored; 2 where the man was colored and the woman white; one where the man was white and the woman colored; in all, 63, or 6 more than in 1891.

The number of marriages where one of the parties was colored and the other white have been as follows in the last eleven years: 1881, 5; 1882, 5; 1883, 1; 1884, 3; 1885, 6; 1886, 4; 1887, 7; 1888, 4; 1889, 4; 1890, 2; 1891, 5; 1892, 3.

The aggregate and average age of the colored persons married in 1892 were as follows:

62 males......Aggregate age, 1,794 years......Average age, 28.93 years. 61 females.....Aggregate age, 1,548 years......verage age, 25.38 years.

The difference in the number of males and females is owing to the fact of the mixed marriages just alluded to. The same difference of course will be observed in the aggregate number given below for the period of thirty-seven years, 1856 to 1892, inclusive. The whole number of marriages of colored persons during that time, either one or both parties being colored, was 1,623; the aggregate and average age of the parties were as follows:

Table X. shows the number and percentage of brides and grooms in each division of age:

Table X. Marriages, 1892. Ages of the Parties.

		AGES OF BRIDES.										ber 8.	Jo
OF GROOMS.	Under 20 years.	20 to 25.	25 to 30.	30 to 35.	35 to 40.	40 to 45.	45 to 50.	50 to 55.	55 to 60.	60 to 65.	65 to 70.	Whole Number of Grooms.	Percentage Grooms.
Under 20 years.	21	7										28	1.76
20 to 25	124	353	67	10	1						• • • •	555	34.86
25 to 30	45	234	185	35	5	1						505	31.72
30 to 35	10	61	82	55	12	2	1					223	14.01
35 to 40	3	23	29	<b>2</b> 8	29	10	1					123	7.73
40 to 45	2	7	8	22	14	6	5	1				65	4.08
45 to 50		4	5	5	7	11	6	2				40	2.51
50 to 55				2	8	6	4	2		1		23	1.44
55 to 60				3	3	3		3	1			13	.82
60 to 65				.1	2	1	2	1	2		1	10	-63
65 to 70					2	1		1		1		5	-31
70 to 75						1		1				2	.13
Whole number of Brides	205	689	376	161	83	42	19	11	3	2	1	1,592	
Percentage of Brides	12.88	43.28	23.62	10.11	5.21	2.64	1.19	.69	.19	•13	.06		100.00

In 658 marriages, or 41.33 per cent. of all, the bride and groom were in the same division of age.

The aggregate and average age of all parties married during the year were as follows:

1,592 malesAggregate a	age,	45,731	yearsAverage	age,	28.72	years.
1,592 femalesAggregate	age,	40,516	$\mathbf{years}\mathbf{Average}$	age,	25,45	years.

The following figures show the statistics for the period of thirty-seven years, 1856 to 1892,; by periods of five years from 1856 to 1890, for the years 1891 and 1892, and for the whole period. For

particulars of each year, see Reports for 1882, page 28, and for 1885, page 24:

		Males.		Fema	ales.
Years.	No.	Aggregate age.	Average age.	No. Aggrega	te age. Average age
<b>1856</b> 1860	3,037	84,831 years	27.93 years	$3\ 034\ 72,971$	years 24.05 years
1861 1865 .	3,152	90,329 years	·28.66 years	3,152 77,353	years 24.54 years
1866-1870	4,428	125,681 years	.28.38 years	4,428109,462	years24.72 years
1871 1875	5,432	155,507 years	28.63 years	5,432135,426	years24.93 years
1876-1880	5.221	152,003 years	.29.11 years	5,221131,646	years25.21 years
1881-1885	5,786	169,588 years	29.31 years	5,786148,515	years25.67 years
1886-1890	6,731	195,871 years	.29.10 years	6,731171,169	years 25.43 years
1891	1,488	46,705 years	.30.02 years	1,488 40,030	years 26.40 years
1892	1,592	45,731 years	.28.72 years	1,592 40,516	years 25.45 years
			-		
37 years 8	6,867	1,066,246 years	.28.92 years	36,864857.088	years 23.25 years

Number of times married. Table XI. shows how many of the marriages in 1892 were the first, second or third marriages of the parties:

Table XI. Marriages, 1892. Number of times Married.

GROOMS.	Nu	Whole Number of			
Number of the Marriage.	First.	Second.	Third.	Fourth.	Grooms.
First	1,257	110	1		1,368
Second	123	84	3	1	211
Third	5	8			13
Whole Number of Brides	1,385	202	4	1	1,592

In 1,257 cases, or 78.96 per cent. of all, it was the first marriage of both parties. The number of grooms married for the first time was 1,368, or 85.93 per cent. of all; of brides, 1,385, or 87.00 per cent.

In 84 cases, or 5.28 per cent. of all, it was the second marriage of both parties.

Marriages of divorced persons. There were, in 1892, 10 marriages where both parties had been divorced; there were 33 other marriages where the man had been divorced, and 63 where the woman had been; in all, 106 marriages, or 6.66 per cent. of all, where one or both of the parties had been divorced. This was 17 more marriages and 0.68 per cent. more than in 1891.

The number and proportion of the marriages of this class in each year since 1883 have been as follows:

Year.	No.	Per ct.	Year.	No.	Per ct.
1883	95	8.07	1888	82	6.08
1884		7.95	1889	97	7.09
1885		6.78	1890	95	6.64
1886	76	6.08	1891	89	5 98
1887	100	7.48	1892	106	6.66

There were 3,184 persons married in 1892 of whom 116, or 3.64 per cent. had been divorced; 43 men and 73 women. These 106 marriages of divorced persons were performed by clergymen of the different denominations, or by civil authority, as follows:

Methodist         .28           Baptist         .23	Episcopalian3  Disciples of Christ2
Congregational11	Evangelical 2
Universalist 8	Justice Supreme Court2
Free Baptist 5	Presbyterian 1
Unitarian 5	Evangelical Advent 1
Christian 5	Latter Day Saints 1
Advent Christian 4	Not specified 5
l l	

Denominational. The 1,592 marriages in 1892 were performed by elergymen of various denominations, or by civil authority, as follows:

#### DENOMINATIONAL.

Roman Catholic648	Hebrew 14
Episcopal206	United Presbyterian 9
Baptist	Evangelical Advent 6
Methodist150	Advent Christian 6
Congregational123	New Jerusalem 5
Free Baptist 44	Disciples of Christ 4
Unitarian 29	Evangelical 3
Universalist	Church of Emanuel 2
Christian 26	Latter Day Saints 1
Lutheran 23	Church of Christ 1
Presbyterian	
Denomination not stated 20	Total
Justice of Supreme Court 18	

Nativity and Residence. There were in 1892, 125 marriages, or 7.85 per cent. of all, in which both parties were natives of Providence. There were besides, 133 marriages where the groom only was born here; and in 235 the bride only. This gives a total of 618 persons married here during the year who were natives of the city; this was 19.41 per cent. of the 3,184 persons married during the year.

In 193 marriages, or 12.12 per cent. of all, both parties were non-residents of the city; this was 4 less marriages and 1.12 per cent. less than in 1891. Of these 193 marriages there were 102 where both parties were from outside the State; in 82 both parties came from Massachusetts; and in 10 others one of the parties was from that State. But the number of marriages of non-residents has decreased since the change in our marriage law, July, 1881.

Marriage Licenses. There were 1,708 marriage licenses issued in 1892 from the office of the City Registrar; this was 116 more than the number of marriages during the year. These two numbers never correspond exactly; see Report for 1886, page 26.

The number of marriage licenses issued in each year since July 1, 1881, is as follows:

1881 (six months) 616	1887
1882	1888
1883	1889
1884	1890
1885	1891 1,607
18 86 1,343	1892

Marriage and Education. The 3,184 persons married in Providence in 1892 were divided as follows, as to sex and nativity:

Born in the United	StatesMales	, 900Females,	, 913Tot	al, 1,813
Born in foreign cou	intriesMales	s, 692Females	, 679 Tot	al, 1,371

The number and proportion who signed the marriage certificate with a mark is shown in the following table. Many certificates are returned with the place for the signature left blank, and it is impossible to know whether this omission is due to inability to write, or to the neglect of the person performing the ceremony in not requiring the signature. These failures to sign are shown in a column by themselves, and the last two columns show the number and proportion of those who signed with a mark, and those who failed to sign, taken together:

Signed	Total In
Whole No. with In each Failed In each	h marks each
1892. Married. a Mark. 100. to sign. 100. s	failures. 100.
Males born in United States 900 22 2.44 10.11.	23 2.55
Females born in United States. 913 21 2.3022.41	43 4.71
Total born in United States1,813 43 2.37231.27	66 3.64
Males born in foreign coun-	
tries 69210214.74 71.01	1 915.75
Females born in foreign coun-	
tries 679 8111.93375.45	11817.38
Total born in foreign coun-	
tries	22716.56
Totals3,1842267.10672.10	293 9.20

There were 43 persons (22 males and 21 females) born in the United States, who signed the marriage certificate with a mark. This was 8 more than in 1891; of these 43 persons, 8, or 18.60 per cent. (3 males and 5 females) were of white American parentage; in 1891 the proportion was 17.14 per cent.; of the other 35 persons, 18 were colored (11 males and 7 females), and 17 of foreign parentage (8 males and 9 females).

There were 183 persons born in foreign countries (102 males and 81 females) who signed with a mark; this was 1 less than in 1891. The whole number of persons signing with a mark in 1892 was 7 more than in the preceding year, and the proportion to the whole number of persons married 0.26 per cent. less.

The following figures show these facts for the period of thirty-three years, 1859 to 1891:

1859-1891.	Whole number Married.		100.
Males born in United States	20,988	****** 889*****	4.24
Females born in United States	21,080	1,192	5.65
	-	-	
Total born in United States	42,068	2,081	4.95
Males born in foreign countries	12 481	3,076	24.64
Females born in foreign countrie	g12,389	4.314	34 82
a common soria in roronga countries			
Total born in foreign countrie	s24.870	7.390	29.71
Totals		.,	

#### DEATHS.

The number of deaths reported in Providence for the year 1892 was 2,964, which was 334 more than in 1891. The number of male decedents was 176 more, of females 158 more; whites 321 more, colored 13 more; natives of United States 152 more, of foreign countries 183 more; of American parentage 96 more and of foreign parentage 236 more; unknown, 1 less. The nativity of 4 decedents and the parent nativity of 2 were not stated.

The population of Providence for July 1, 1892, was estimated at 138,000.\* In the following table the number in each class is calculated by assuming that it formed the same percentage of the total population that it did in 1890 as shown by figures sent through the courtesy of the Superintendent of the Eleventh Census:

`	Population, 1892.	Deaths, 1892.	To population one death in	Deaths in each 1,000 of population.
Total population	138,000	2,964		21.48
Males	76,391	1.506	50 72	19.71
Females	61,609	1.458	2.12	23.67
Whites	133,763	2 821	47.41	21.08
Colored	4,237	143		33.75
Born in United States.	95,841	2,008	47.73	20.95
Born in foreign countri	ies. 42 159	952	44.28	22.58
American parentage	54,579	1,179	46.29	21.60
Foreign parentage	83,421	1,783		21.37

The rate of mortality in 1892 was 2.00 more than in the preceding year. During the thirty-seven years from 1855 to 1891 inclusive, there were 62,274 deaths in Providence, giving an average annual death-rate of one to each 50.48 of the population, or 19.81 deaths in each 1,000 living. The death-rate in 1892 was therefore 1.67 more than the average for the thirty-six preceding years.

<sup>\*</sup>A census taken in January. 1893, gave the population of the city as 148,944.

TABLE XII. DEATHS, 1892.

Sex, Condition, Color, Nativity, Parentage, Locality and Season.

1892.	January.	February.	March.	April.	May.	June	July.	August	September.	October	November	December	YEAR.
Whole number of Deaths.	347	240	229	221	234	231	324	256	246	199	220	217	2,964
Sex.— Males Females	163 184	115 125				119 112			121 125	99			
Condition.— Married. Single. Widows. Widowers. Divorced. Unknown.	125 63 20	70 116 35 18 1	82 106 31 8 2	97 23 14	105 34 10	131 26 9	75 217 21 9 2	155 21 11	76 132 24 11 3	98 24 12	108 29 16	25	946 1,487 356 150 20 5
Colored	328 19	231 9	214 15	206 15		220 11	312 12		234 12	193 6	211 9	208 9	2,821 143
Nativity.— United States. Ireland. England and Scotland. Germany. British America. French Canada. Portugal & Western Isl. Sweden. Norway. Italy. Other Countries. Unknown.	219 76 26 4 6 7 2 3 	160 50 12 2 6 3 2 1	144 52 14 4 2 5 2 1 	138 48 15 3 7 1  3 5	157 47 13 2 5 2 3 2 	4	242 46 17 3 2 4  1	188 46 17 1 3 1	170 55 7 4 4 3 1	142 33 13  3 1 1 2  2	148 46 11 3 4 3 2 2 2	144 31 19 3 4 5 1 2	2,008 568 178 33 50 41 14 16 1 27 24 4
Parentage.— American Irish. English, Scotch & Welsh German French Canadian Portuguese Swedish. Norwegian. Italian Other Foreign Unknown Foreign	148 118 37 8 5 9 6 4 2 5 4 1	102 88 19 4 6 4 1 5 2 5 4	92 85 18 5 4 8 3 1 1 7 3 2	86 75 26 5 4 3 3 1	104 78 22 2 2 6 6 2  6 3 3	92 79 22 5 3 12 1 2 1 6 5 2	109 116 33 8 5 18 5 8 1 19 14 8	99 86 28 7 5 8 5 3 1 5 5 4	93 100 22 6 9 6 1 2  4 2 1	80 71 25 4 2 6 3 2	82 83 22 3 6 6 7 4 	92 63 29 4 2 6 2 4  8 6 1	1,179 1,042 303 61 53 87 43 38 8 68 59 21
Locality, Wards.—													
I	36 31 32 29 39 34 34 31 27 17	14 26 31 16 22 16 19 13 23 36 24	29 22 26 14 19 17 25 11 19 24 23	19 16 27 10 13 15 16 22 27 35 21	18 17 27 9 16 23 16 22 23 23 40	22 20 21 11 22 22 12 17 27 26 31	33 20 38 8 26 29 24 18 31 58 39	21 18 27 16 15 23 15 21 27 34 39	19 12 28 12 17 9 20 19 25 46 39	10 19 36 8 18 14 9 13 16 32 24	21 17 25 7 20 12 22 24 11 25 36	20 17 27 11 14 18 16 24 20 26 24	262 235 345 151 241 232 228 235 276 402 357
Totals	347	240	229	221	234	231	324	256	246	199	220	217	2,964.

Season. The largest monthly total of deaths in 1892 was 347, in January. This was the largest number of deaths ever recorded in one month, except January, 1890, when the total was 384.

The following figures show the number of deaths in each quarter of each of the last six years; also the aggregate and average number of deaths in each quarter of the year, in the total for thirty-seven years, from 1855 to 1891, inclusive:

						37 years, 1855-1891.
	1892.	1891.	1890.	1889.	1888.	1887. Aggregate. Average.
January-March	816	607	864	638	717	63115,314 414
April-June	686	618	635	584	582	59314,133 382
July-September	826	665	754	717	694	71417,584
October-Dec	636	740	624	571	615	67415,243 412
				· .		
Totals	2,964	2,630	2,877	2,510	2,608	2,61262,2741,683

The largest number of deaths was in the third quarter of the year and the smallest in the fourth quarter.

The following shows the number and percentage of deaths in Providence in each quarter of the year, in the aggregate for 31 years, from 1855 to 1885, inclusive; for the period of five years, 1886 to 1890, inclusive, and for the years 1891 and 1892:

	JanM	arch.	April-June.	July-Sept.	OctDec.
	Deaths.	Per ct.	Deaths. Per ct.	Deaths. Per ct.	Deaths. Per ct.
1855-1885	11,283	24.17	.10,59822.70	13,34228.58	11,45924.55
1886-1890	3,424	26.42	. 2,91722.50	3,57727.60	3,04423.48
1891	607	23.08	61823.50	66525.28	74028.14
1892	816	27.53	68623.14	82627.87	63621.46

Sex and Parentage. The sex of the decedents, by parentage, in Providence, is shown in the following table, in periods of five

years each, from 1861 to 1890, inclusive, and for the years 1891 and 1892:

American Parentage.	Foreign Parentage.
Males. Females. Total.	Males. Females. Total
1861-1865 1,401 1,474 2,875	1,401 2,796
1866-1870 1,407 1,499 2,906	1,387 1,332 2,719
1871-1875 1,978 2,158 4,136	2,150 2,192 4,345
1876-1880 2,179 2,354 4,538	2,593 2,772 5,368
<b>1</b> 881–1885 2,380 2,635 5,01 <b>5</b>	3,011 3,086 6,09
1886-1890 2,673 2,807 5,480	3,696
1891 531 552 1,083	799 748 1,54
1892 596 583 1,179	908 875 1,788
32 years 13,145 14,062 27,207	15,94516,18632,13

The parentage of two male decedents was unknown.

Civil Condition. The 2,964 decedents in 1892 and the 2,630 decedents in 1891 were divided as follows as to condition and parentage:

Civil	American	Parentage	e.	Foreign	Parentage.
condition.	1891.	1892.	Total.	1891.	1892. Total.
Married	330	311	. 641	503	6351,138
Single	540	611	1,151	824	8751,699
Widows	138	175	. 313	151	181 332
Widowers	66	65	131	58	85 143
Divorced	9	14	. 23	6	6 12
Unknown		3	. 3	5	1 6
Totals 1	1,0831	,179	2,262	1,547	1,7833,330

Of the 2,262 decedents of American parentage in the two years 1,151, or 50.88 per cent., were single, and 1,111, or 49.12 per cent., were married, or had been.

Of the 3,330 decedents of foreign parentage, 1,699, or 51.02 per cent., were single, and 1,631, or 49.98 per cent., were married, or had been; of ten, the civil condition was not stated (3 of American, 6 of foreign and 1 of unknown parentage).

Color. The white decedents of 1892 were 2,821 in number, 1,040 of American, 1,179 of foreign parentage and 2 unknown.

There were 139 colored decedents, which was 9 more than in 1891; 4 of these were of foreign parentage.

Nativity and Parentage. The nativity and parentage of the decedents is given in Table XII. The following figures show what difference there is between the two classes:

1892.	Nativity.	Parenta	ge
United States	2,008	American1	179
Ireland	568	Irish1	,04%
England and Scotland	178	English and Scotch	308
Germany	33	German	61
British America	50	British-American	58
" (French)	41	French-Canadian	8
Portugal and Western Island	14	Portuguese	48
Sweden	16	Swedish	38
Norway····	1	Norwegian	. 8
Italy	27	Italian	68
Other countries	24	Other foreign	59
Unknown	4	Unknown foreign	21
		Unknown	6
Totals	2,964		,96

The record of the decedents under these two classifications is important, as may be seen by the difference between the numbers as given above. The number of decedents born in the United States was 2,008, or 67.75 per cent. of all, but the number of American parentage was only 1,179, or 39.77 per cent. of all. The decedents of foreign birth numbered 952, or 32.12 per cent. of all; those of foreign parentage 1,783, or 60.16 per cent.

The decedents born in other foreign countries than those named in the table were as follows: Born in Russia 7, Armenia 5, France 4, Austria 2, and one each in Asiatic Turkey, Bermuda, Denmark, Holland, Poland and Switzerland.

There were 59 decedents of "other foreign" parentage, as follows: Russian 25, French 8, Armenian 5, Danish 3, Polish 3, Swiss 3, Austrian 2, Dutch 2, Roumanian 2, Belgian, South American, St. Helena, Syrian, Turkish and West Indian one each.

Locality. The following figures show the number of decedents, by parentage, in each ward of the city, in 1891 and 1892:

	American.		Fore	ign.		Amer	Foreign.		
Wards.	1891.	1892.	1891.	1892.	Wards.	1891.	1892.	1891.	1892
I	121	90	154	172	VII	123	142	60	8
II	106	124	106.	111	VIII	127	133	96	10
III	69	86	278.	259	1X	96	87	190	18
IV	86	83	56.	68	x	66	81	252	32
V	112	112	109.	129	Public In-				
vi	99	104	116.	128	stitutions	78	137	130	21
							-		
Wh	ole city.					1.0831	.1791	.547	1.78

The decedents in public institutions in each of the last five years have been as follows. It will be seen from the table that the term *Public Institutions*, for this purpose, includes not only hospitals and other public institutions, but also hotels, rivers, ponds, the streets, etc., where persons happen to die away from their usual residences:

	1892.	1891.	1890.	1889.	1888
Rhode Island Hospital	134	91	96	84	8
Butler Hospital	14	15	16	17	1
Homeopathic Hospital	14	8	7	11	
St. Joseph's Hospital	21	0	0	0	
Dexter Asylum	17	24	19	19	1
Home for Aged Men	2	1	5	2	
Home for Aged Women	5	9	1	8	
Home for Aged Colored Wome	n 0	1		0	
Children's Home					
St. Elizabeth Home		0	3	1	
st. Maria's Home					
Shelter for Colored Children					
Rhode Island Nursery					
St. Aloysius Asylum					
small Pox Hospital					
st. Vincent de Paul Asylum					
Working Men's Home					
R. C. Convent				-	
state Home and School					
Lying-In Hospital					
seaman's Bethel					
Hotels					
Police Station					
Bays, Rivers and Ponds					
others					
Jnknown					

Aggregate and Average Age, and Season. The following table (XIII.) shows the number of deaths, and the aggregate and average age of the decedents, in Providence, according to parentage, in each month, and in the whole year 1892:

Table XIII. Deaths, 1892.

Aggregate and Average Age by Parentage.

	Ameri	ican Par	entage.	Forei	ign Pare	entage.	Total	Americ Foreign	
MONTHS, 1892.	Number of Deaths.	Age. Average Age. Age.		Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.
January	148	7,302	49.34	199	8,324	41.83	347	15,626	45.03
February	102	4,310	42.25	138	4,569	33.11	240	8,879	37.00
March	92	3,010	32.72	137	5,288	38.59	229	8,298	36.24
April	86	3,214	37.37	135	4,812	35.64	221	8,026	<b>3</b> 6.32
May	104	4,335	41.68	130	4,558	35.06	234	8,893	38.00
June	92	2,642	28.71	139	4,291	30.87	231	6,933	30.01
July	109	2,397	21.99	215	4,690	21.81	324	7,087	21.87
August	99	2,413	24.37	157	4,225	26.91	256	6,638	25.93
September	93	2,820	30.32	153	5,148	33.65	246	7,968	32.39
October	80	3 064	38.30	119	3,732	31.36	199	6,796	34.15
November	83	3,357	40.45	137	4,793	34.77	220	8,150	37.04
December	92	4,077	44.32	125	3,909	31.27	217	7,986	36.80
Totals	1,180	42,941	36.39	1,784	58,339	32.70	2,964	101,280	34.17

The average age of decedents of American parentage was 3.69 years greater than that of decedents of foreign parentage. The average age of all decedents in 1892 was 0.74 years more than in 1891; of decedents of American parentage 0.49 less, and of foreign parentage 1.69 more.

The following table, XIV., gives the aggregate and average age of the decedents, by months and by parentage, in the aggregate for the thirty-seven years from 1856 to 1892, inclusive:

TABLE XIV. DEATHS, THIRTY-SEVEN YEARS, 1856-1892.

Recapitulation. Aggregate and Average Age by Parentage.

	Americ	an Pare	ntage.	Foreig	n Paren	tage.		Americai Foreign.	n and
THIRTY-SEVEN YEARS, 1856-1892.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.			Number of Deaths.	Aggregate Age.	Average Age.
January	2,775	102,628	36.98	2,927	87,057	29.74	5,702	189,685	33.27
February	2,377	85,652	36.03	2,603	72,729	27.94	4,980	158,381	31.80
March	2,539	92,239	36.33	2,696	76,620	28.42	5,235	168,859	32.26
April	2,479	88,496	35.70	2,746	80,691	29.38	5,225	169,187	32.38
May	2,392	89,075	37.24	2,586	77,333	29.90	4,978	166,408	33.43
June	2,035	72,765	35.76	2,373	66,206	27.90	4,408	138,971	31.53
July	2,690	69,520	25.84	3,545	74,668	21.06	6,235	144,188	23.13
August	2,869	80,994	28,23	3,552	76,493	21.54	6,421	157,487	24.53
September	2,375	71,164	29.96	3,040	72,552	23.87	5,415	143,716	26.54
October	2,310	79,273	34.32	2,822	73,606	26.08	5,132	152,879	29.79
November	2,303	80,691	35.04	2,717	75,891	27.93	5,020	156,582	31.19
December	2,631	97,274	36.97	2,865	83,669	29.20	5,496	180,943	32.92
Totals	29,775	1,009,771	33.91	34,472	917,515	26.62	64,247	1,927,286	30.00

The average age of American decedents during this period of thirty-seven years was 7.29 years greater than that of the foreign class.

In the reports for 1882, 1885 and 1886 more extended comparisons are made between the two classes of decedents, in the matter of age.

Percentage of decedents at different ages. The next table (XV.) shows the number and percentage of the decedents, by parentage, in the different divisions of ages in 1892, and also in the aggregate for thirty-seven years, from 1856 to 1892, inclusive:

Table XV. Deaths, 1892.

				Amer Parer 18	itage,	Fore Parer 189	itage,	To Ame & For 189	rican reign,	37 Years, 1856-1892.	
			AGES.	Number of Deaths,	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.
Un	der	1 yea	r	291	24.66	348	19.51	639	21.56	12,481	19.43
1.8	and	unde	r 2	52	4.41	78	4.37	130	4.38	4,769	7.42
2	46	66	ō	57	4.83	74	4.15	131	4.42	5,136	7.99
	To	tal uı	nder 5	400	33.90	500	28.03	900	30.36	22,386	34.84
5 :	and	unde	r 10	38	3.22	40	2.24	78	2.63	2,900	4.51
10	6.6	6.6	15	28	2.37	36	2.02	64	2.16	1,252	1.95
15	4.6	6.6	20	34	2.88	41	2.30	75	2.53	2,081	3.24
20	6 6	6.6	30	60	5.08	197	11.04	257	8.67	6,394	9.95
30	6.6	6.6	40	67	5.68	221	12.39	288	9.73	5,882	9.16
40	6 6	4.6	50	77	6.53	203	11.38	280	9.44	5,160	8.03
<b>5</b> 0	66	85	60	75	6.36	208	11.66	283	9.55	5,148	8.01
60	6.6	6.6	70	137	11.61	192	10.76	329	11.10	5,233	8.15
70	66	46	80	140	11.86	105	5.88	245	8.26	4,773	7.43
80	46	44	90	110	9.32	34	1.91	144	4.86	2,553	3.97
90	and	over		14	1.19	7	.39	21	.71	485	-76
	То	tal	••••	1,180	100.00	1,784	100.00	2,964	100.00	64,247	100.00

The following table shows the ratio of deaths under one year to births according to parentage, in periods of five years each from

1856 to 1890, and for 1891 and 1892, and for the whole period of thirty-seven years from 1856 to 1892:

RATIO OF DEATHS.

	AMERICAN F	ARENTAGE.	FOREIGN PARENTAGE.				
Periods.		Ratio of Deaths under 1 year to Births.	Births.	under	Ratio of Deaths under 1 year to Births.		
1856-1860	3,621 431				12.43		
1861-1865	3,056 414	13.54	4,147	593	14.30		
1866-1870	4,149 511	12.31	4,681	595	12.72		
1871-1875	5,415 787	14.53	6,323	970	15.34		
1876-1880	6,102 810	13.27	6,848	1,013	14.79		
1881-1885	. 7,091 995	14.03	7,308	1,099	15.04		
1886-1890	7,3581,136	15.71	8,284	1,327	16.01		
1891	1,698 251	14.78	2,178	325	14.92		
1892	1,601 291	• • • • • • 18.17	2,251	348	15.46		
1856-1892	40,0915,626	14.33	46,727	6,855	14.67		

The following table shows the percentage of decedents in Providence, under 5 years of age, according to parentage, in periods of five years each, from 1856 to 1890; also in 1891 and 1892, and in the whole period of thirty-seven years, from 1856 to 1892, inclusive:

Table XVI.

Percentage under 5 years according to Parentage.

	age,	can Par 37 year 56-1892.		Foreign 37 ye	Paren ars, 18 1892.	itage,	American and Foreign Parentage, 37 years, 1856-1892.			
PERIODS.	Total Deaths.	Deaths under 5 years.	Percentage under 5 years.	Total Deaths.	Deaths under 5 years.	Percentage under 5 years.	Total Deaths.	Deaths under 5 years.	Percentage under 5 years.	
1856–1860	2,567	856	33.35	2,340	1,218	52.05	4,907	2,074	42.27	
1861-1865	2,875	882	30.68	2,796	1,342	48.00	5,671	2,224	39.22	
1866-1870	2,906	804	27.67	2,719	1,128	41.49	5,625	1,932	34.35	
1871-1875	4,136	1,297	31.36	4,342	1,788	41.18	8,478	3,085	36.39	
1876-1880	4,533	1,547	34.13	5,365	2,145	39.98	9,898	3,692	37.30	
1881-1885	5,015	1,613	32.16	6,097	1,859	30.49	11,112	3,472	31.25	
1886-1890	5,480	1,880	34.31	7,482	2,299	30.73	12,962	4,179	32.24	
1891	1,083	365	33.70	1,547	463	29.93	2,630	828	31.48	
1892	1,180	400	33.90	1,784	500	28.03	2,964	900	30.36	
37 years,	29,775	9,644	32.39	34.472	12,742	36.96	64,247	22,386	34.84	

These figures show that the percentage of decedents of American parentage, under 5 years, has varied very little throughout the entire period; while the proportion of foreign decedents of that age has steadily and greatly diminished.

During the 15 years, from 1840 to 1854 inclusive, of all the decedents in the city, 43.56 in each 100 were under 5 years of age; from 1856 to 1880 inclusive, of all the decedents 37.55 in each 100 were under 5 years; from 1881 to 1891, of all decedents, 31.75 in each 100 were under 5 years, showing a progressive decrease in the mortality at this age.

In the 53 years, 1840 to 1892 inclusive, there were 77,392 decedents in Providence, of whom 28,121, or 36.57 per cent., were under 5 years of age. The number and percentage in each division of age under 5 years were as follows:

Whole number of decedents in 53 years77,392
Under one year of age14,98619.36 per cent. of all-
One and under two years 6,552 8.47 per cent. of all.
Two and under five years 6,583 8.51 per cent. of all.
Total under five years

The following table (XVII.) shows the number and percentage, by parentage, of the decedents in each division of ages, in the aggregate of decedents in the 37 years, from 1856 to 1892 inclusive:

Table XVII. Deaths, 1856-1892.

Number and percentage at different ages, according to parentage.

	American age, 37 1856-	Parent- years, 1892.	Foreign age, 37 1856-	years,	American and Foreign Parentage 37 years, 1856-1892.		
AGES.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	
Under 1 year	5,626	18.90	6,855	19.88	12,481	19.43	
1 and under 2	1,891	6.35	2,878	8.35	4,769	7.42	
2 and under 5	2,127	7.14	3,009	8.73	5,136	7.99	
Total under 5	9,644	32.39	12,742	36.96	22,386	34.84	
5 and under 10	1,350	4.53	1,550	4.50	2,900	4.51	
10 and under 15	597	2.00	655	1.90	1,252	1.95	
15 and under 20	884	2.97	1,197	3.47	2,081	3.24	
20 and under 30	2,328	7.82	4,066	11.80	6,394	9.95	
30 and under 40	2,372	7.97	3,510	10.18	5,882	9.16	
40 and under 50	2,227	7.48	2,933	8.51	5,160	8.03	
50 and under 60	2,394	8.04	2,754	7.99	5,148	8.01	
60 and under 70	2,778	9.33	2,455	7.12	5,233	8.15	
70 and under 80	3,042	10.22	1,731	5.02	4,773	7.43	
80 and under 90	1,847	6.20	706	2.05	2,553	3.97	
90 and over	312	1.05	173	.50	485	.76	
Totals	29,775	100.00	34,472	100.00	64,247	100.00	

The difference in the percentages of the two classes of decedents at different periods of life may be seen by the following statement:

American parentage, under 50 years 19,402or	65.16	per	cent.
American parentage, 50 years and over	34.84	4.6	6.6
Foreign parentage, under 50 years	77.32	6.6	6.6
Foreign parentage, 50 years and over	22.68	6.6	6.6

Deaths by age, parentage and season. The next table (XVIII.) shows the number of decedents, by parentage, in Providence, in each division of ages, in each month and each quarter of the year 1892. The table shows considerable difference in the proportion of decedents, by parentage, in different seasons of the year.

Table XVIII. Deaths by age and parentage, in each month and quarter, 1892.

	DIVISIONAL PERIODS OF LIFE.																
D	IVISIONS OF	មព្		1				og	9 PER	Se cross	60	.0.	80.	.06	100	<b>D</b> .	ses.
	THE YEAR.	Under 1 Year.	1 to 2.	2 to 5.	5 to 10.	10 to 1	15 to 2	20 to 3	30 to 4	40 to 5	50 to 6	60 to 7	70 to 8	80 to 9	90 to 1	100 and Over.	All Ages.
	January February March	19 20 22	4 1 7	7 10 5	1 3 2	5 2 2	2 2 1	4 5 5	8 5 8	11 5 10	10 5 6	27 12 10	29 14 11	19 14 3	24		148 102 92
	1st Quarter.	61	12	22	6	9	5	14	21	26	21	49	54	36	6		342
AGE.	April May June	19 20 29	1 1 6	5 3	3 6 3	2 3 2	1 6 6	5 6 7	12 5 2	4 7 6	8 3 7	6 13 9	10 16 4	9 11 7	1 4 1		86 104 92
PARENTAGE.	2d Quarter.	68	8	11	12	7	13	18	19	17	18	28	30	27	6		282
JAN PAB	July August September	49 45 27	9 5 7	4 4 3	6 2 4	1 2 4	2 2 2	3 4 4	3 5 5	6 5 8	3 4 6	13 5 7	4 11 10	6 5 6			109 99 93
AMERICAN	3d Quarter.	121	21	11	12	7	6	11	13	19	13	25	25	17	• •		301
An	October November December	13 14 13	2 4 5	4 6 3	5 2 1	1 2 2	6 3 1	6 5 6	2 3 9	5 3 7	10 6 7	11 13 11	7 9 15	8 12 10	2		80 82 92
	4th Quarter	40	11	13	8	5	10	17	14	15	23	35	31	30	2		254
	Whole Year	290	52	57	38	28	34	60	67	77	75	137	140	110	14		1,179
_	January February March	25 21 14	6 4	7 8 7	3 4	2 6 2	3 5 1	21 11 16	21 15 20	23 14 15	22 29 21	37 13 19	21 5 10	9 1 3	.1		199 138 137
	1st Quarter.	60	16	22	9	10	9	48	56	52	72	69	36	13	2		474
sá.	April May June	18 18 30	7 5 4	4 6 9	1 3 3	2 4 1	1 3 7	21 13 22	18 22 10	21 16 12	19 11 17	12 18 11	7 9 6	4 2 4			135 130 138
TAG	2d Quarter.	66	16	19	7	7	11	56	50	49	47	41	22	10	2		403
PARENTAGE.	July August September	84 45 32	11 13 7	7 6 5	4 2 2	6 1 2	4 3 4	19 12 15	20 22 17	28 15 17		10 14 22	5 8 9	1 2 3	 i	1	215 157 153
FOREIGN	3d Quarter.	161	31	18	8	9	11	46	59	60	46	46	22	6	1	1	525
FOH	October November December	27 13 21	6 6 3	1 9 5	5	28	1 5 4	15 17 15	17 21 18	17 13 11	10 21 12	9 16 11	10 6 9	1 3 1	i		119 137 125
	4th Quarter	61	15	15	16	10	10	47	56	41	43	36	25	5	1		381
	Whole Year Unknown Parentage	348 (Nov.)1	78	74	40	36 	41	197	221	<b>2</b> 02 (Jun.)1	208	192	105	34	6	1	1,783 2
	otal Amer- an & Foreign	639	130	131	78	64	75	257	288	280	283	329	245	144	20	1	2,964

The whole number of American decedents was 1,179; of decedents of foreign parentage, 1,783. The percentage of each class in the different quarters of the year was as follows:

	1st Quarter.	2d Quarter.	3d Quarter.	4th Quarter.
American parentage	29.00	23.91	25,53	21.54
Foreign parentage	26.58	22.60	29.45	21.36

Table XVIII. shows the relative mortality at different ages and seasons. Of the 301 American decedents in the third quarter, 121, or 40.20 per cent., were less than a year old. The whole number of American decedents under one year was 290; the 121 who died in the third quarter were 41.58 per cent. of this number.

Of decedents of foreign parentage, the number in the third quarter was 525; of these, 161, or 30.67 per cent., were less than one year old; the whole number of foreign decedents of this age was 348; the 161 dying in the third quarter of the year were 46.26 per cent. of the whole number.

Causes of Death. Table XIX. gives the number of deaths in Providence in 1892 from each cause stated, and in each division of ages; also the number of decedents from each cause, according to sex and parentage; the percentage of deaths from each cause, of the whole number from known causes, and also the number of deaths from each cause to 100,000 of the population.

Season and Disease. Table XX. shows the number of deaths from some of the most prominent causes, in each month and quarter of the year 1892, showing the comparative mortality from different diseases at different seasons.

Table XIX. Deaths, 1892. Disease, sex, age, parentage, percentage from known causes, and ratio per 100,000 of population.

100,000 tion.	Deaths per slugod to	
s from whole mwonn	Hatio of Death each cause to Mo, from l causes.	<u> </u>
GE.	Total.	- 4400 0 8 8 7 7 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
PARENTAGE	Foreign.	480000005-44400 - 400000888800000
PAI	American	<u>∞.</u> σ.
	J9VO 35 06	: st = : : : : : : : : : : : : : : : : : :
	,08 of 08	
	,08 of 07	0 4 5 5 F F F F F F F F F F F F F F F F F
IFE.	.07 of 08	4
DIVISIONAL PERIODS OF LIFE	.00 to 60.	
DS G	.08 ot 04	SON SOSSALVHES HEDOMACH
ERIO	\$0 to 40.	Ø\$ØØ\$ : ₩₽ : ₩ : ₩ : ₩ : Ø\$# Ø 4 ₩ : Ø\$Ø₽
T P	20 to 30.	
NA	.02 of dI	::::::::::::::::::::::::::::::::::
ISIC	.čI ot 0I	: D : : St = : : : : : : : : : : : : : : : : : :
	.01 of č	
	2 to 5.	
	1 to 2.	
	Under I year,	
	Total.	- 440 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
SEX.	Females.	05-75-1-50 1-4 1-4 1007-1- 105828210212
	Males.	
	CAUSES OF DEATH.	A becesses.  A cridents (various).  Burns and seads.  Burns and seads.  Borowning.  Falls.  Poisoning.  Alroholism.  Aneurism.  Appendicitis.  Appendicitis.  Alle Duct., obstruction of Brit.  Bills Duct., obstruction of Brit.  Brit., obstruction of Brit.  Brit., or Stomach.  Cancer (various).  Cancer (various).

Table XIX. Deaths, 1892. Disease, sex, age, parentage, percentage from known causes, and ratio per 100,000 of population.

.по;оот	reathsed Singoq to	######################################
s from mwond	Hatio of Death each cause to No. from causes.	
GE	Total.	<u> </u>
PARENTAGE.	Foreign.	######################################
PAI	American	<u> </u>
	• 19 VO 3 09	
	.08 ot 08	
별	.08 of 07	
LIE	*07 03 08	
DIVISIONAL PERIODS OF LIFE.	.00 to 60.	
RODS	40 to 50.	
Per	.04 of 08	
AL	20 to 80.	co ·
NO	15 to 20.	
ISI	.df of 0f	:::::::::::::::::::::::::::::::::::::::
Úrv	.01 o1 d	
	2 to 5.	<u>\$\frac{\partial}{\partial}\tag{\partial}\ta</u>
	I year.	- 73 : 78 % 14 % . % 14 %
-	Under	:F:: : : : : : : : : : : : : : : : : :
	Total.	194 Reg. 86044
SEX.	Females.	480 84-0408r-5.4505 -4208-400 -11
	Males.	88-1888 8214 614 418 44 44 44 51 14 14 14 14 14 14 14 14 14 14 14 14 14
	CAUSES OF DEATH.	Child Birth, convulsions Cholera Infantum. Colic

Table XIX. Deaths, 1892. Disease, sex, age, parentage, percentage from known causes, and ratio per 100,000 of population.

100,000 ation.	req satae Singoq to	6848444824348835846848585858584858485
known whole trom	Ratio of Death each cause to No. from causes,	
GE.	Total.	1000 100 100 100 100 100 100 100 100 10
PARENTAGE.	Foreign.	14-14-11:03-44:11-08-04:1-04:1-08:05-09-08:11-08
PA1	American	
	- 19VO & 00	
	.08 of 08	10 10 10 10 10 10 10 10 10 10 10 10 10 1
ಣೆ	.08 of 07	
CIE	.07 ot 0a	
DIVISIONAL PERIODS OF LIFE	.09 of 03	
ODS	40 to 50.	
PERI	.04 of 08	
AL.	.08 of 02	
ION	15 to 20.	:=::::::::::::::::::::::::::::::::::::
ISI	.dI of 01	: : : : : : : : : :
)IV	.01 of d	
	2 to 5.	
	1 to 2.	
	Under 1 year.	
	Total.	
SEX.	Females.	
	Males	
	CAUSES OF DEATH,	in Remittent in Remittent in Typhoid in Typhoid Gangrene General Paralysis Heart, Diseases of Hydrocephalus Lalundice Kidneys, Diseases of Laryngtils Laucocythæmia Leucocythæmia

Table XIX. Deaths, 1892. Disease, sex, age, parentage, percentage from known causes, and ratio per 100,000 of population.

100,000 tion.	Deaths per	4-10-17
whole known	Ratio of Death each cause to No. from causes.	
GE.	Total.	<u>848-02-02-02-084-688-28-1-00-00-1-889-1</u>
PARENTAGE.	Foreign.	24747-64 :: 884-87-804-48
PA	American	100
	1. TAVO 38 08	::::::::::::::::::::::::::::::::::::::
	.09 of 08	260 O H44E0 H 67
	.08 of 07	
DIVISIONAL PERIODS OF LIFE	.07 od 08	
OF	.00 to 60.	
SOC	.0d of 04	66.000
ERIC	.04 of 08	
LE	.08 of 02	27 8 7 8 8
NA	15 to 20.	::::::::::::::::::::::::::::::::::::::
Sro	10 to 15.	: H : : : : : : : : : : : : : : : : : :
IVI	5 to 10.	
D	to 5.	
	1 to 2.	os ·∞ — os ·π — — — os
	Under 1 year.	200 ro4 H 4-004
	Total.	#48-50-00-00-05-0-05-0-05-0-05-0-05-0-05-
SEX.	Females.	
	Males.	2 1 1 1 1 2 1 2 2 2 2 2 2 2 2 2 2 2 2 2
	CAUSES OF DEATH.	Lungs, Diseases of Malfornation Malfornation Marsanus Majaranus Measles. Meningitis, Cerebro-Spinal Mearlysis Odd Are. Gangrene. Paralysis. Peritonitis Perenancy, extra-uterine. Prograto, Diseases of. Prograto, Diseases of. Prograto, Diseases of. Rachitis Rheumatism. Scarlet Fever

Table XIX. Deaths, 1892. Disease, sex, age, parentage, percentage from known causes, and ratio per 100,000 of population.

Sex. D	Males. Females. Total. Ivear. I to 2. I to 2.	Septicarmia	1,506 1,458 2,964 639 130 131
DIVISIONAL PERIODS OF LIFE	10 to 10. 10 to 15. 15 to 20. 20 to 30.		78 64 75 257
ERIO	30 to 40.		288
D8 01	.00 00 05	\$0. 50 com	280 283
F LIF	*02 O1 09	© 1 1 1 0 1 1 1 0 1 1 1 1 1 1 1 1 1 1 1	3 -329
8	.08 ot 07		245
	.08 of 08		144 21
PA	American		1,180
PARENTAGE	Foreign.	\$\$\$\$P\$\$\$4 : \(\pi\)\$\$14\(\pi\)\$\$1 : \(\pi\)\$\$1 : \(\pi\)\$ : \(\pi	1,784
i ii i	Total.	840000000000000000000000000000000000000	2,964
known be whole	Ratio of Deat each causet No. from causes.	<u>6444588628854845488888888888888888888888</u>	100.00
100,000 . noits	Deaths per	4.824.444 7.448.86.67 8.86.44 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.75 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.75.75.45 1.7	21.48

Table XX. Deaths, 1892. Season and Disease.

9 :	2854552558486888852845844855°
Whole Year.	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~
Fourth Quarter.	\$\$\$4\$\$240 :H444H2:T050400834000
December.	ರ್ವಾಹದ್ದರ್ಧ : ಸ್ಥಾತ್ರೀಯ : ಸ್ವಾಪ್ತ್ : - ಸ್ವಾಪ್ತ್ರ : - ಸ್ವಾಪ್ತ್ : - ಸ್ವಾಪ್ತ : - ಸ್ವಾಪ್ತ್ : - ಸ್ವಾಪ್ತ್ : - ಸ್ವಾಪ್ತ್ : - ಸ್ವಾಪ್ತ್ : - ಸ್ವಾಪ್ತ : - ಸ್ವಾಪ್ತ : - ಸ್ವಾಪ್ತ : - ಸ್ವಾಪ್ತ್ : - ಸ್ವಾಪ್ತ : - ಸ್ವಾಪ್ತ : - ಸ್ಟ್ರಾಪ್ಟ್ : - ಸ್ಟ್ರಾಪ್ಟ್ : - ಸ್ಟ್ರಾಪ್ಟ್ : - ಸ್ಟ್ರಾಪ್ಟ್ : - ಸ್ಟಿಪ್ಟ್ : - ಸ್ಟ್ರಾಪ್ಟ್ : - ಸ್ಟ್ಟ್ಟ್ಟ್ಟ್ಟ್ಟ್ಟ್ಟ್ಟ್ಟ್ಟ್ಟ್ಟ್ಟ್ಟ್ಟ್ಟ್ಟ
November.	Short Salar
October.	ÖÖ ü ü Ü ü 4 ü 4 ₽ 4 Ø ₽ . 2 8 8 . 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
Third Quarter.	48888808808807878800488
September	Ferriousisseries
August.	
July.	Racolica6820 40F245 - 4-E 24F4 13
Second Quarter.	<u> </u>
June.	0040m : # :
.vam	Arredas
April.	
First Quarter.	88222221111112451-51-x471 : 884 × 1 € 1 € 1 € 1 € 1 € 1 € 1 € 1 € 1 € 1
Матер.	ASTERON REPUSAAND ST SEC - OUR SIL
February.	<u> </u>
January.	# 1 - 66 - 10 - 10 - 10 - 10 - 10 - 10 -
CAUSES OF DEATH.	Accidents (all kinds).  Apoplexy Brain, Disasses of Brain, Disasses of Cancer. Child Birth Cholera Infantum Cholera Infantum Convulsions. Covup Coupp Debility Distribus Diphtheria Diphthe

Table XX.—Concluded. Deaths, 1892. Season and Disease.

lle r.	0824588 155 155 165 165 165 165 165 165 165 165	2,964
Whole Year.		
Fourth Quarter.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	636
Decemper.	20.01.02.1.1.1.2.2.2.2.2.2.2.2.2.2.2.2.2	217
November.	φυ	220
October.	2001-1201-1 1001-00 100 NO	199
Third Quarter.	8470881 .:: 24108418418	828
September.		246
August.	© 000 00 00 00 00 00 00 00 00 00 00 00 0	256
July.	20 H 65 K 4 H H 65 65 65 65 65 65	324
Second Quarter.	:: :::::::::::::::::::::::::::::::::::	989
June	4010081 :000 :01-4-01 :00	231
May.	Umwastarua . 4	234
April.	Ф : : : : : : : : : : : : : : : : : :	221
First Quarter.	8 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	816
March.	ου : 84-ου : σο : -ου Ε	559
February.	## ## ## ## ## ## ## ## ## ## ## ## ##	240
January.	11.00-26.24.44.4	347
CAUSES OF DEATH.	Old Age Paralysis Pertionitis Pertinonitis Puthisis Preumonia Congestion of Lungs. Congestion of Lungs. Scarlet Fever Scarlet Fever Scarlet Fever Scarlet Fever Storach, Diseases of. Storide Tuberculosis Tuberculosis Whooping Cough. Whooping Cough. Unknown.	Totals

# NOTES ON THE CAUSES OF DEATH.

### DURING THE YEAR 1892.

These notes are in explanation of the facts given in tables XIX. and XX.; there are also given additional statistics, and comparisons with the statistics for the thirty seven years, 1856 to 1892.

The percentage of deaths from the different causes has always been reckoned on the number of deaths from known causes.

The number of deaths in Providence in 1892 was 2,964; of these there were 14 cases, or 0.47 per cent., where no cause of death could be determined or recorded; the number from known causes, therefore, and on which all the percentages are reckoned was 2,950.

In the thirty-seven years, 1856 to 1892 inclusive, there were 64,247 deaths in Providence; 676, or 1.05 per cent., for which no cause was given, the number from known causes being 63,571. In the 53 years, 1840 to 1892 inclusive, there were 77,392 deaths; of 1,678 or 2.17 per cent., the cause was not stated; the number of deaths from known causes in this period being 75,714.

Abscess. The number of deaths from abscess in 1892 was 7, the same number as in 1891. The location of the abscesses was as follows: 3 pelvic; 1 psoas; 1 hip; 1 ovary; 1 face and brain.

All deaths from known causes 2,950.
From abscess, 7, or 0.24 per cent. of all.
Male, 1; females, 6.
American parentage, 3; foreign parentage, 4.

For the 37 years, 1856 to 1892, the figures are:

All deaths from known causes, 63,571.

Deaths from abscess, 180, or 0.28 per cent. of all.

Males, 89; females, 91.

American parentage, 79; foreign parentage, 101.

Accidents. There were 121 deaths from accidents in 1892, or 42 more than in the preceding year. There were 10 deaths by asphyxia; 9 from burns and scalds; 13 by drowning; 25 from falls; 5 from poisoning, and 11 by railroad accidents; 48 from various other accidents. The exact manner of all these accidents is not always described; so far as specified they are as follows:

Asphyxia. One of the deaths from asphyxia was due to food in the larynx. The others were due to illuminating gas.

Burns and Scalds. Matches 5; hair catching from gas 1.

Falls. From ladder 2; from bed, from sofa, from roof, from wagon, from window, into hold of schooner and during syncope, 1 each.

Poisoning. By over-dose of opium, by over-dose of laudanum, by Paris-green and by illuminating gas, 1 each.

Various Accidents. There were 48 accidents of various kinds which caused death as follows: Fracture of femur 3; of hip 5; of skull 5; of leg 1; of fore-arm 1; of tibia and scapula 1; of spine 1; of pelvis 1; hit by falling telegraph pole 1; puncture of bladder 1; caught in machinery 3; horse-car accident 1; cable-car accident 1; blasting accident 1; buried in sand 5; gun-shot wound of head 1; pistol-shot wound of hand 1; hit by runaway horse 2; powder explosion 1; concussion of brain 2; explosion of steam cylinder 2; wound of thigh, scrotum, perineum and rectum 1; thrown from team 1; kicked by horse 1; hit by falling derrick 1; injury to thumb 1; run over by low gear 1; elevator accident 1; thrown from horse 1.

The general statistics of deaths from accidents are as follows:

Males, 91; females, 30; total 121, or 4.11 per cent. of all. American parentage, 40; foreign parentage, 81.

Aneurism. There were 3 deaths from aneurism in 1892, the same as in 1891, and 0.10 per cent. of all deaths from known causes.

2 males; 1 female; total 3, or 0.10 per cent. of all. American parentage, 1; foreign parentage, 2. In the 37 years, 1856 to 1892, there were:

Males, 45; females, 15; total, 60, or 0.09 per cent. of all. American parentage, 24; foreign parentage, 36.

Apoplexy. This was given as the cause of 98 deaths in 1892; 9 more than in 1891.

Males, 54; females, 44; total, 98, or 3.32 per cent. of all.

American parentage, 39; foreign parentage, 59.

The number of deaths from this couse has increased greatly during the past thirty years, as is shown by the following table, which gives the number in each period of five years from 1856 to 1890, inclusive; also for 1891 and 1892, and for the 37 years, 1856 to 1892, inclusive.

	Deaths per 100,000 of			Deaths per 100,000 of
Deat	ns. Population.	1	eaths.	Population.
1856-1860 87	33	1881-1885	394	68
1861-1865 86	32	1886-1890	491	- 79
1866-1870 134	43	1891	89	66
1871–1875 177	42	1892	98	71
1876-1880 240	47 ·	_		_
37 vears			1796	56

Total deaths from known causes in 37 years, 63,571.

From apoplexy: Males, 854; females, 942; total, 1796, or 2.83 per cent. of all.

American parentage, 1,095; foreign parentage, 701.

Bones, Diseases of. The one case of bone disease, was of the ankle joint.

Brain, Diseases of. The number of deaths reported from diseases of the brain in 1892 was 79, or 8 more than in 1891. Of these 79 deaths 53 were called inflammation; 8, congestion: 6, softening; 2, tumor, and one each, atrophy and abscess;

Males, 39; females, 40; total, 79, or 2.68 per cent. of all. American parentage, 43; foreign parentage, 36. For the 37 years, 1856 to 1892 the statistics are:

Males, 1,235; females, 1,060; total, 2,295, or 3.61 per cent. of all. American parentage, 1,209; foreign parentage, 1,086.

Bronchitis was returned as the cause of 148 deaths in 1892; this was 31 more than in the preceding year.

Males, 64; females, 84, total 148, or 5.02 per cent. of all.

For the 37 years, 1356 to 1892, the decedents were:

Males, 692; females, 832; total, 1,524, or 2.40 per cent. of all. American parentage, 661; foreign parentage, 863.

For full particulars, see the table on the two following pages:

BRONCHITIS IN PROVIDENCE, THIRTY-SEVEN YEARS, 1856-1892.

Sex; parentage; season; age; whole number and ratio to known causes.

-siatoT	1,524	692 832	661 863	89111 8800 852 852 852 852 852 852 852 852 852 852 852
.268I	148	64	90	883110014100
.1681	117	670	48	ಬರಿಂಬ್ಯಾಣಕಾರ್ಣ ಹಣ್ಣಾ ಅಭ
*0681	133	66	72	0.04F055r050000
.6881	107	56	65	084854548104
.8881	3.30	38 488	57	555120000000000
.788I	3.18	47 36	29	15-45-2000000000
.9881	82	255	38	∞о <u>∺ы</u> шии4иио4∞
.3881	3.57	977	43 43	@F#014@@@@F&@
.4881	2.78	888	85.42	F-070F-70-4-4-0-70-0-4
.6881	57.	23.88	42.88	7272700724601H8000
.2881	60	22	22,88	F-1040000000000
.1881	53	88	38.7	<u> </u>
.0881	56	229	85.55	%00004H040H0
.9781	34	23.53	18	04704880 SH .48
.8781	2.38	18	19	; 
.7781	4.5	22.83	1935	∞
.9781	46	8.8	28	L0447888L4L40
.8781	33	17	128	00000
.4781	1.68	12	9.74	
.8781	1.05	801		_ :: \( \text{index} \) : \( \text{index} \)
.2721	10.1	431	2-6	
.1781	17.	10	9 11	40000000000
.0781	17.1	<b>&amp;</b> @	<b>c.</b> ∞	erenene :- :-eren
*6981	1.21	411	96	: : : : : : : : : : : : : : : : : : :
.8981	10.92	. 9	€5.00	
.7981	106	≃ ເ	4.9	S4 :::
.9981	0.39		- 63	
1861 to	8.0	12 21	25	
1856 to	26 0.55	16	55	4 44%000-000-
BRONCHITIS.	Whole number	SEX: Males Females	PARENTAGE: American Foreign	SEASON: January February March March June June July Agrith Cottober October December

Bronchitis in Providence, Thirty-Seven Years, 1856-1892.—Concluded.

Sex; parentage; season; age; whole number and ratio to known causes.

Totals.	10 10 10 10 10 10 10 10 10 10 10 10 10 1	521 397 206 400	694 354 476
.2681	0000188889985748	28827	330
.1981	88801	08 15 4 8 15 8	42 21 54
.06SI	: : : : : : : : : : : : : : : : : : :	# <del>\$</del>	3000
.6881	230118818819248	24.00 10.00	22.22
*8881	E STEWNHOWSHOWS :	35 24 11 16	23 19
.7881	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	35 0 24 24	39 29 29
.9881		22.23.23	15 26
.6881	<u> </u>	23 11 17	20.03
.488I	801100111111111111111111111111111111111	120181	18 18
.588I		15 17 19 19	144
.2881		118	30
.1881	£ £ € € € € € € € € € € € € € € € € € €	41 81 82 81	20 16 17
.0881	24411 H HOUSE	10 10 10 10 10 10	25.
.6781	□ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □	7.00	10
.8781	0004H	10,822	27 10 10
.7781	4 mm : H44mm	167 9	111181
.9781		12882	122
.č78I	1	110	13
.478I	044 : :	9 8 4 1	10
.8781	1 10 4 4 4 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1		999
.2781	1000H	704864	∞ \$5 €
.1781		∞ 20 – r∪	ರಾಜ್
.0781	33 :03 :	ಎಸುಟ4	x 410
·698I	रान : : नकानक्ष्यं	5-403	040
.8981	w . H	⊗=∞4	ಯಬಹ
.7381	ණ දැක් දැක් දැක් දැක් දැක් දැක් දැක් දැක්	9 : 1	9-6
.9981		o: ⊢:	: = ::
1861 to 1865.	000000000000000000000000000000000000000	21.094	17 9
1856 to 1860.	000	∞24r0	55 8 9 .
RONCHITIS.	Under 1 year.  I and under 2.  I in (10.  I	quarter I quarter th quarter	four monthsad four months.
BF	AGE:: 200 100 100 100 100 100 100 100 100 100	First que Second Third que Fourth	First fo Second Third f

The lowest annual percentage of deaths from bronchitis in the 37 years, was 0.34 in 1862; the highest 5.02 in 1892, and next to the highest 4.65 in 1890. There is a close connection between this disease and pneumonia; and while the statistics of each are given separately, the facts are compared under the head of pneumonia.

Cancer. The number of deaths from this cause in 1892 was 77, or 4 less than in 1891. Of these, 15 were cancer of the stomach; 13 of the uterus and 11 of the breast. The 38 others, so far as specified, were, cancer of the abdomen 5, of the liver 4, of the rectum 3, of the intestines, throat, lip, and pelvis 2 each, and of the omentum, of the bladder, of the omentum and bladder, of the face, ventriculi, of the shoulder, of the pancreas, of the kidney, of the mouth, of the cheek, of the gall-duct, of the urethra, 1 each.

Males, 21; females, 56; total, 77, or 2.61 per cent. of all. American parentage, 41; foreign parentage, 36.

For the 37 years, 1856 to 1892, the statistics are:

Males, 469; females, 1,276; total, 1,745, or 2.74 per cent. of all. American parentage, 1,025; foreign parentage, 720.

The following table shows the sex, parentage, whole number and percentage of the decedents from cancer, by periods of five years, from 1856 to 1890, inclusive; for 1891 and 1892; and for the 37 years, 1856 to 1892:

DECEDENTS FROM CANCER, 37 YEARS, 1856-1892.

	Sex.		Parenta	ge.	Whole	
Years.	Males.	Females.	American.	Foreign.	Number. Perce	entage
856-1860	20	. 61	56	25	81	1.
861-1865	31	. 79	80	30	110	1.9
866-1870	36	. 95	100	31	131	2.3
1871-1875	54	. 152	128	78	206	2.
1876-1880	84	. 217	161	140	301	3.
1881-1885	92	. 267	211	148	359	3.
1886-1890	110	. 289	209	190	399	3.
1891	21	. 60	39	42	81	3.
1892	21	. 56	41	36		2.
		. —				

The percentage of males among these decedents was 26.88; of females, 73.12; of American parentage, 58.74; of foreign parent-

age, 41.26. Full particulars of each year may be found in the report for 1888, page 55.

Childbirth. There were 27 deaths from this cause in 1892, which was 7 more than in 1891. Twelve deaths were from puerperal fever, and 4 from puerperal convulsions. The decedents were 8 of American and 19 of foreign parentage; total 27, or 0.92 per cent. of all.

In the 37 years, 1856 to 1892, the decedents were:

666, or 1.05 per cent. of all. American parentage, 250; foreign parentage, 416.

Cholera Infantum. The number of deaths from this cause in 1892 was 151, or 33 more than in the year preceding.

Males, 83; females, 68; total, 151, or 5.12 per cent. of all.

American parentage, 65; foreign parentage, 86.

The following table shows the sex, parentage and percentage of the decedents from cholera infantum in 37 years, 1856 to 1892:

	Sex.		Parenta	ge.	Whole	
	Males. 1	Females. A	merican.	Foreign.	Number.	Percentage
856 to 1860	138	119	94	163	257	
861 to 1865	139	140	92	187	279	
866 to 1870	175	149	124	200	324	
871 to 1875	278	297	217	358	575	6.
876 to 1880	243	230	181	292	473	4.
881 to 1885	294	263	218	339	557	
886 to 1890	324	305	279	350	629	4.
			43			
			65			

The 1,313 decedents of American parentage were 39.04 per cent. of all; the 2,050 foreign decedents 60.96 per cent.

Until the year 1883, the above statistics of cholera infantum were given by single years, so that by reference to the report of that and succeeding years, the full statistics of this disease may be found for each separate year.

Cholera Morbus. There were 10 deaths from cholera morbus in 1892; 5 more than in 1891.

Males, 7; females, 3; total, 10, or 0.34 per cent. of all. American parentage, 5; foreign parentage, 5. In the 37 years, 1856 to 1892, there were 231 deaths from cholera morbus, or 0.36 per cent. of all.

Convulsions. There were 53 deaths caused by convulsions in 1892, which was 11 more than in the preceding year. The decedents were:

Males, 35; females, 18; total, 53, or 1.80 per cent. of all. American parentage, 27; foreign parentage, 26.

For the 37 years the statistics were:

Males, 869; females, 764; total, 1,633, or 2.57 per cent. of all. American parentage, 689; foreign parentage, 944.

In addition to the above, convulsions were given as the secondary and immediate cause of death in 31 cases in connection with the following diseases: Bright's disease and albuminuria 5, meningitis and teething 4 each, indigestion 3, cholera infantum, cerebral congestion and diarrhœa, 2 each, stomatitis, gastro-intestinal irritation, diabetes, pneumonia, fall, bronchitis, cholera morbus, congestion of the lungs and malaria, 1 each.

Croup. The number of deaths from this cause in 1892 was 34, or 18 more than in 1891. The number and per cent. in each quarter of the year were:

Number. Percent.	Number. Per cent.
First quarter1544.12	Third quarter 2 5.88
Second quarter 3 8.82	Fourth quarter1441.18

#### The decedents were:

Males, 20; females, 14; total, 34, or 1.15 per cent. of all. American parentage, 19; foreign parentage, 15.

The general statistics for the 37 years, 1856-1892, were:

Whole number, 1,156; annual average, 31
Males, 588; females, 568: total, 1,156, or 1.82 per cent. of all.
American parentage, 470; foreign parentage, 686.

The next table shows the statistics of croup in Providence, during 37 years, from 1856 to 1892, inclusive, showing the sex, color and parentage of the decedents; the season and locality of the deaths, and the percentage of the deaths from croup, of all the deaths from known causes:

DEATHS FROM CROUP IN PROVIDENCE, THIRTY-SEVEN YEARS, 1856-1892.

[3toT]	1,156	588 568	470 686	621 628 628 628 628 628 638 638 638 638 638 638 638 638 638 63	141 125 128 118 118 119 119 128 138 141 158
.2681	34	30	15		SSB SS
.1681	16.0.61	51	421		
.0881	1.08	22	13		40-00-4 704 -
·6881	33	119	10	8888 8 aadaa	00 10000000 00 ·
.8881	1.04	4.65	4.60	**************************************	
.7881	39	18	41 25	00000000000000000000000000000000000000	1 18987-1-18
.9881	32	121	13	0000-00-1-00-4	∞=× • ∞×=∞ ≈ = : :
.8881	1.30	12	55.50	: : : : : : : : : : : : : : : : : : :	H :00000000400 H
.488I	33	16	13	; 5-44000 :00==4000	
.6881	35	1789	13	F-0500 0005 007070-44	о ;« :ниимою н
.288I	1.43	14	45	; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ;	0 :=05= :1045-0 =
1881	49	25.83	32	F-00000000-40F-F-	: ::::::::::::::::::::::::::::::::::::
.0881	29	21.	222		g :4Lg :Hprop
.6781	2.13	27	22.2	orv∞466641-61	∞ € ∞ − m ∞ ≥ ∞
.8781	38 1.93	18	18	47000 HH 61000	20-00-0505-45-H 00
1.7781	2.07	18	14 36		F-014 - 01 - 01 - 01 - 01 - 01 - 01 - 01
.9781	65 3.48	83 83	3827	16°55° 23° 12°65°	απα4ωαωφαπα :
.6781	2.95	28.23	37	10000 : 11000000	0 4 442
748T	1.48	100	15	.:: .::	4 H : SH475-4 H
.678I	122	15	120	\$1000000000000000000000000000000000000	4444 .0050
.2781	1.77	13	9 21 8		4 :F-034HH070 - H
.1781	2.83	18	20	: ::::::::::::::::::::::::::::::::::::	онинфии4F : :
.0781	2,16	16	12	44444 - 40040	. :
*698T	1.53	10	133	тан	: : : : : : : : : : : : : : : : : : :
.8881	1.01	100	415-		8 T 8 8 8 T
.788I	19	10	10		4600-100 : 60
.9981	2.27	122	122	; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ;	
1861 to 1865	162 2.86	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~	63	22222422410	22-84588 · · · · · · · · · · · · · · · · · ·
1856 to 1860	123	69	725	81123500044153	₹ £ £ £ £ £ £ £ £ £ £ £ £ £ £ £ £ £ £ £
CROUP.	Whole number Percentage	SEX: Males Females	PARENTAGE: American Foreign	SEASON. January January March. May June July September. October. October. December.	Ward I. Ward II.  "" III "" V. "" VI "" VII "" VIII "" VIII "" VIII "" VIII "" VIII "" VIII

The ward totals on this table do not include the figures after 1887, as those are reckoned by the new division of wards; and all before that are for the old wards of the city.

The number and proportion in each quarter of the year, for this period of years, were as follows:

Dea	ths. Per cent.	Deaths. Per cer	ıt.
First quarter41	3 35.73	Third quarter 129 10.	.38
Second quarter18	88 16.26	Fourth quarter435 37	.63

These figures show that 848 deaths, or 73.36 per cent. of all the deaths from croup, occurred in the colder season of the year, from October to March, inclusive.

Comparisons between the statistics of croup and diphtheria are given under the head of the latter cause.

Debility. This was the cause assigned for 87 deaths in 1892, of which 65 were debility from premature birth. Of the 22 others, 14 were less than a year old:

Males, 53; females, 34; total, 87, or 2.96 per cent. of all. American parentage, 42; foreign parentage, 45.

For the thirty-seven years, 1856 to 1892.

Whole number, 1,152; annual average, 31.

Males, 604; females, 548; total, 1,152, or 1.81 per cent. of all.

American parentage, 516; foreign parentage, 636.

Diarrhea and Dysentery. The number of deaths from diarrhea in 1892, was 55; from dysentery, 30; in all, 85; or 2.89 per cent. of all; 26 more deaths and 0.63 per cent. more than in 1891.

These and other diseases classed under the head of diarrheal are included in the following table, showing the number, sex, parentage and percentage from each cause in 1892:

Males	Females.	American.	Foreign.	Total. Per cent.
Acute Diarrhœa 17.	29	16	30	46 1.56
Chronic Diarrhoea 4.	5	4	5	931
Dysentery 16	14	4	26	30 1.02
Cholera Infantum 83.	68	65	86	151 5.12
Cholera Morbus 7.	3	5	5	10
Enteritis 17.	15	14	18	32 1.08
Enteritis, Gastro 26	20	14	32	46 1.56
Enteritis, Colo 7	13	7	13	2068
Enteritis Muco	1		1	1
Totals,	167	129	215	344 11.67

The whole number of deaths from diarrheal diseases was 113 more than in 1891. The decedents from these causes in the thirty-seven years, 1856 to 1892 were:

Whole number, 6,677; annual average, 180.

Males, 3,342; females, 3,335; total, 6,677, or 10.50 per cent. of all.

American parentage, 2,722; foreign parentage, 3,955.

The proportions of American and foreign parentage were as follows:

	American. Foreign.
1892 166.6 foreign to 100 American	; or 37.50 and 62.50 in each 10
36 years previous144.2 foreign to 100 American	; or 40.94 and 59.06 in each 100

Of all decedents in 1892, the proportions were 151.1 foreign to 100 American; or 60.19 foreign and 39.81 American in each 100 decedents.

The following table shows the number of deaths from each diarrheal disease, and the percentage from all these diseases, in different periods of time, from 1840 to 1892, inclusive:

Ratio to all.						Cholera	Cholera	
known causes.	l'otal. k	eritis.	ry. Ent	Dysente	Jiarrhœa.	Morbus.	Infantum.	
	1,870	102		796 .	349	63	····560	1840-1854
12.06	. 573	23		127	159	14	250	1855-1859
13.27	. 703	46		189 .	154	17	297	1860-1864
12.15	665	37		187	125	35	281	1865-1869
12.04	. 931	97		91.	185	26	532	1870-1874
9.18	. 883	104		77	169	24	509	<b>1</b> 875–1879
9.78	1,068	164		74	220	39	571	1880-1884
9.0	1,107	220		121	175	53	538	<b>1</b> 885–1890.
8.88	231	49		23 .	36	5	118	1891.
	344	98		30 .	55	10	151	1892.
		_	_					
		98	-	30 .	55	10	151	

The Deaths from Asiatic cholera are not included in the above figures; they were 158 in 1849; 159 in 1854; and 14 in 1866; total, 331.

Diphtheria. There were 39 deaths from this cause in 1892, or 9 less than in the preceding year, and less than in any year since

1885. The number and proportion in each quarter of the year were as follows:

	Deaths.	Per cent.	Deaths. Per cent.
First quarter	10	25.64	Third quarter 6 15.39
Second quarter	10	25.64	Fourth quarter

## The decedents were:

Males, 20; females, 19; total, 39, or 1.32 per cent. of all. American parentage 12; foreign parentage 27.

The first deaths reported from diphtheria in Providence were in 1858. During the thirty-five years, 1858 to 1892, there were 2,050 decedents from this disease.

Decedents from all known causes in 35 years, 61,643.

From diphtheria, 2,050; annual average, 58.

Males, 1,014; females, 1,036; total, 2,050, or 3.33 per cent. of all.

American parentage, 956; foreign parentage, 1,094.

The following table shows the whole number, the percentage, season and locality of the deaths from diphtheria, also the sex and parentage of the decedents in the thirty-five years, 1858 to 1892, inclusive. The statistics are given for the three years, 1858 to 1860; the five years, 1861 to 1865; the five years, 1866 to 1870, and for each year since 1870:

DEATHS FROM DIPHTHERIA IN PROVIDENCE, THIRTY-FIVE YEARS, 1858-1892.

Totals.	3.33	1,014	956 1,094	198 150 1149 1178 1178 178 250 250	253 106 1158 1143 2555 2555 2555 2555 2555 2555 2555 25
1892.	1.32	19	122	まちょうなものできますら	する41000437-00
1891.	1.83	22.	32	4400040034070	
1890.	3.29	43	47	HH-01-40001-80	800000041007
1889.	3.90	253	42	01 01-01 02-44 00 00 00 01	70 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
1888.	3.77	24 CC	50	0.041.004804.51 <b>0</b> .85	941147-08
1887.	1.19	55	78	11 8 87 - 00 4 0 2 E E S 8	8 :32 :11 :22 :23
1886.	4.17	46	500	45-70000000475-4	<u> </u>
1885.	1.71	18	16	© m m m m m ≥ 4 m m m m ∞	
1884.	2.60	8%	36	œ <b>™⊕⊔</b> 4∺∞ым∞	<u> </u>
1883.	2.33	8.28	22.22	<b>⊕</b> თ∢•••∞∞∞∞∞∞4	4200000000000
1882.	2.15	88	33.8	rosener ronner E	000000
1881.	116	55	57	27-78 211 27-11 24 4 4 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	50 H 455 4 0 55 85 85
1880.	2.95	88	25.55	1-04040001010101010	0830 LVII-80
1879.	5.30	46	468	Orena Orena Sala	F484851088
1878.	246 12.47	136	105	2018210 o 2018888	8444911488 44491488
1877.	295	146	129	4250822°2888	582428188 58248188
1876.	5.95	10 to	6,88	10 4707000510000000000000000000000000000000	C00040005000
1871 to 1875.	105	52.52	88	5255-455-1000	24.001.018.4
1866 to 1870	0.79	23.	21.2		∞ +4 co co co co →4 co ≥+ :
1861 to 1865.	138	99	83	10 10 10 10 10 10 10 10 10 10 10 10 10 1	221232123
1858 to 1860.	1.66	8.4	42.83	::::::::::::::::::::::::::::::::::::::	u4∞€10000 · · ·
DIPHTHERIA.	Whole number	SEX; Males Females	PARENTAGE: American Foreign	SEASON: January March March May July July September October December	Locality: I I I I I I I I I I I I I I I I I I I

The ward totals in this table do not include the figures after 1887; as those are reckoned by the new division of wards, and all before that are for the old ward totals.

The subject of "diphtheria" has been commented upon in several previous reports; see "City Registrar's Report" for the years 1860; 1870; 1877 to 1880, inclusive; and 1883.

The following table illustrates the subject of season in connection with the three important causes of death, diphtheria, scarlet fever and croup; showing the number and percentage of deaths in Providence, from each cause, in each month and quarter of the year. The figures for diphtheria cover thirty-five years, 1858 to 1892, inclusive, for scarlet fever, the thirty-eight years, from 1855 to 1892, inclusive, and for croup, the thirty-seven years, from 1856 to 1892, inclusive. Until the year 1884, the figures for croup included the year 1855; but in the reports since that year, the figures concerning that disease for 1855 are omitted from this comparative table, that the whole number may correspond with the total from croup, as given in the table on page 66, which covers the period from 1856 to the present time:

	1858-	HERIA, -1891, ears.	SCARLET 1855- 37 ye		CROUP, 1856-1890, 36 years.		
Months.	Number of Deaths.		Number of Deaths.	of Per cent.		Per cent.	
January	193	9.42	313	12.19	159	13.76	
February	150	7.32	270	10.51	125	10.81	
March	161	7.85	230	8.96	129	11.16	
1st quarter	504	24.59	813	31.66	413	35.73	
April	138	6.73	226	8.80	85	7.35	
May	149	7.27	225	8.76	64	5.54	
June	120	5.85	188	7.32	39	3.37	
2d quarter	407	19.85	639	24.88	188	16.26	
July	101	4.93	134	5.22	39	3.37	
August	104	5.07	110	4.28	18	1.56	
September	178	8.68	122	4.75	63	5.45	
3d quarter	383	18.68	366	14.25	120	10.38	
October	271	13.22	208	8.10	121	10.47	
November	250	12.20	234	9.11	160	13.84	
December	235	11.46	308	12.00	. 154	13.32	
4th quarter	756	36.88	750	29.21	435	37.63	
Totals	2,050	100.00	2,568	100.00	1,156	100.00	

Dividing the year into periods of four months, we find the following numbers and percentages:

SEASONS.	Дірнт	HERIA.	SCARLET	FEVER.	CROUP.		
	Number.	Per cent.	Number. Per cent.		Number.	Per cent.	
1st four months	642	31.32	1,039	40.46	498	43.08	
2d four months	474	23.12	657	25.58	160	13.84	
3d four months	934	45.56	872	33.96	498	43.08	

Dividing the year into periods of six months, we have:

SEASONS.	DIPHT	HERIA	SCARLET	r Fever.	CROUP.		
	Number.	Per cent.	Number.	Per cent.	Number.	Per cent.	
1st six months	911	44.44	1,452	56.54	601	51.99	
2d six months	1,139	55.56	1,116	43.46	<b>5</b> 55	48.01	

Dropsy was the only cause given for 7 deaths in 1892; 3 more than in 1891.

Males, 4; females, 3; total, 7, or 0.24 per cent. of all. American parentage, 6; foreign parentage, 1.

Enteritis. There were 98 deaths from enteritis in 1892, including 20 from colo-enteritis, and 46 from gastro-enteritis.

Males, 50; females, 48; total, 98, or 3.32 per cent. of all. American parentage, 35; foreign parentage, 63.

The number of deaths from enteritis in the 37 years, 1856 to 1892, was 875, or 1.38 per cent. of all from known causes.

Erysipelas. This was the cause of 7 deaths in 1892, the same as in 1891.

Males, 5; females, 2; total, 7, or 0.24 per cent. of all. American parentage, 2; foreign parentage, 5. In the 37 years, 1856 to 1892, the decedents were:

Males, 148; females, 157; total, 305, or 0.48 per cent. of all. American parentage, 153; foreign parentage, 152.

Fever, Malarial. In 1869, two deaths, and in 1880, one death were reported as due to malarial fever. These were counted in with typhoid fever, but since 1880 the record of malarial fevers has been kept separate from typhoid. Malarial disease originated in Providence in 1880 for the first time in fifty years.

The following table shows the deaths from all forms of malarial fevers since 1880:

YEARS.	Males.	Females.	American.	Foreign.	Whole Number.	Percent-
1880	1			1	1	0.05
1881	1	2	2	1	3	0.14
1882	3	2	3	2	5	0.22
1883	4	2	2	4	6	0.26
1884	5	5	6	4	10	0.45
1885	5	4	2	7	9	0.42
1886	7	8	4	11	15	0.64
1887	18	10	10	18	28	1.07
1888	8	11	4 .	15	19	0.73
1889	9	15	4	20	24	0.96
1890	12	11	8	15	23	0.80
1891	3	5	2	6	8	0.30
1892	5	6	4	7	11	0.37
13 years	81	81	51	111	162	0.51

Fever, Typhoid. The number of deaths from this cause in 1892 was 51, which was 11 less than in 1891.

Males, 35; females, 16; total, 51, or 1.73 per cent. of all. American parentage, 17; foreign parentage, 34.

For the 37 years, 1856 to 1892, the statistics were:

Whole number, 1,883; annual average, 51.

Males, 998; females, 885; total, 1,883, or 2.96 per cent. of all.

American parentage, 907; foreign parentage, 976.

In the 25 years previous to 1881, deaths were occasionally reported from "fever," and "bilious fever," and about 30 in all as "typhus fever." There was also, in 1869, one death reported as "intermittent fever," and one "malarial fever;" and in 1880 one death from "malarial fever." These were all included with typhoid fever.

The following table shows the statistics of typhoid fever, in Providence, for the five years, 1856 to 1860, inclusive, the five years, 1861 to 1865, and for each year since 1865. The figures show the whole number, the percentage, season, and locality of the deaths, and the sex, color, parentage and ages of the decedents:

DEATHS FROM TYPHOID FEVER IN PROVIDENCE, THIRTY-SIX YEARS, 1856-1892.

Sex; parentage; color; season; age; locality; whole number and ratio to known causes.

Totals.	1883 2.96	998 885	907 976	88	127 822 108 103 145 232 244 233 251 251	304 289 456 834
1892.	1.73	35	17.	Н	@rvs/wrs/48005+4	170
.1891.	2.37	255	2.4	03	4070-0000000000	110
.0681	39	24	15	භ	4000001110000000	13
*688I	2.37	18	388	70	70000-01-0400go	2112
.8881	3.95	50	73%	20	: ::::::::::::::::::::::::::::::::::::	21108
.7881	39	19	23.00		\$55-400x-555	51451
.8881	2.25	338	32	_	# : www : - wow = wo	80 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
1885.	2.04	83.83	18	:	<b>∞</b> 4884∞840€∞−	9 113 14
.4881	50.00	88	35	c/s	7070HH48H48E94	13 13 13
1883.	128	772	74	4	ರೂ ಕ್ಷಣ್ಣ ಅಭಾರತಿಯಾಗಿ	86 98 98 98 98
.2881	140	66	83	60	4000000 400000	9 112 13 106
.1881	38	820	22.22	GS	000-000000+0000	100
1880,	553	38.33	24 29	Т	81-8454L88L08	171
*6281	98	19	38	_	-100-1000-10010-	16.53
.8781	2.38	28	202	ಣ		83000
.7781	6.65	88	22	35	00 00 H H 45 - 00 50 00 00	15226
.8781	33	11,	19	63	10-10-01-01-01-01-01-01-01-01-01-01-01-0	<u> </u>
.d78I	3.75	33.38	27	63	87100 :: 100 ses	5,4200
*\$48I	3.06	33	88	Ţ-a	בקהמי בממטרב	∞,°2±88
1873.	3.58	888	20 23	9	4000 : 400 40 EL	230
.2781	68	988	23	9	8 : 68-117888918	10 00 to 00
.1781	288	82.88	88	ಣ	およるますのようものののの	12-12
.0781	52	26	255	CS.	8 :44HHN4708F	66 29 29
*698I	38.3	132	123	:		70 C T T
*8981	24 22 22 2	13	16	:		10 - 10 TC
.7881	3.49	813	23	63	: 	4450
*998I	46	88	17 29	ţ-0	H4H8H886000F0	166
1861 to 1865	3.89	95	88	12	20000000000000000000000000000000000000	8258
1856 10 1860	2.97	73	30 SE	4	811 0 4 87 1 1 8 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	2822
TYPHOID FEVER.	Whole number	SEX: Males Females	PARENTAGE: American Foreign	Colored	SEASON: January February February March May June July August September September November November	First quarter Second quarter Third quarter. Fourth quarter.

DEATHS FROM TYPHOID FEVER IN PROVIDENCE, THIRTY-SIX YEARS, 1856-1892.—CONCLUDED.

causes.
known
to
ratio
and
number
whole
locality;
age;
season;
color;
parentage;
Sex;

Totals.	412 405 1056	10 17 17 11 16 11 14 14 14 14 14 15 16 17 17 17 18 18 18 18 18 18 18 18 18 18 18 18 18	171 73 73 73 73 73 73 73 74 74 75 75 75 75 75 75 75 75 75 75 75 75 75
.268I	18 13 20	:	80000004140
.1881	2148	H : 445-8800000	100000000000000000000000000000000000000
.0681	യ്യല	:- :44@rossu	0000 4 70 00 00 00 str
*6881	24 10 25	::Lww018214-188	-101010101010-1 -10101010101010-1
.888I	113		85-11473
.7881	2554		
*9881	3000	.H .470005-400H	1101201180
.788F	1138	::	4 : 2012 2000
.4881	2112		21-22-22-22
.6881	423.4		22.000000000000000000000000000000000000
.2881	1155	: 1000 880 880 00 1 : :	01000000000000000000000000000000000000
.1881	12 14	: HHUNHOND 4 ON HHH	4 . 24 4 0 00 00 05 00
1880	10 16 27	H :: www.05-w4w :	œೞೞ४488₽₽Ğ
.6781	923	1 :0%4E0HH4%	
.8781	4600	:	
.7781	122		10-00-10-00-00 10-00-10-10-10-00
.0781	27		&1-04r088r0000
.8781	9889	18200082484	<u> </u>
.478I	01 00 02 00 00 00 00 00 00 00 00 00 00 00	HHL00x40x0xch	4-500000000
.8781	111	:-420025E0405-:-	F-10:00F-0:10F-40
.2781	212 40	::: 5,0000000000000000000000000000000000	- 20 Transmar
.1781	2272	H 9708475-00H H :	0.0014100∞0
.0781	34		: 
.e381	100		
.8881	16	ø : : : : : : : : : : : : : : : : : : :	F-0505-1005-405-1
.7881	100	L : 52 4 L O L	704707070HW:
*9981	26.23	. 400407070004	9101904
1861 to 1865	488 127	11233327777	£4464 <b>8</b> 4
1856 10 1860	32	H400001088810000	183343313
TYPHOID 1	First 4 months Second 4 months.	AGE: Under I year. I and under 2 2 4 10 11 11 11 11 11 11 11 11 11 11 11 11	Ward II  Ward II  WILL  WARD  WARD

\*I7 were in R. I. Hospital. \*\*1 in R. I. Hospital. †14 were in R. I. Hospital. †6 in R. I. Hospital. \$4 in R. I. Hospital. ¶3 in R. I. Hospital.

. he ward totals in this table do not include the figures after 1887, as those were reckoned by the new division of wards; and all before that were for the old wards of the city.

Percentage. The 1,883 deaths in 37 years were 2.96 per cent. of all deaths from known causes. The lowest rate during that period was 1.36 per cent. in 1890; the highest, 6.27 per cent. in 1882.

Sex. The decedents were 998 males, and 885 females; a proportion of 112.8 males to each 100 females; or 53.00 males and 47.00 females in each 100.

Parentage. There were 907 decedents of American, and 976 of foreign parentage; 107.6 foreign to each 100 American, or 51.83 foreign and 48.17 American, in each 100.

Season. The deaths were at different seasons of the year, as follows:

	Deaths.	Per cent.	Deaths.	Per cent.
1st quarter	304	16.14	3d quarter456	24.22
2d quarter	289	15.35	4th quarter834	44.29

Dividing the year into three parts, we have:

D	eaths.	Per cent.
First four months	412	21.88
Second four months	405	21.51
Third four months	1,066	56.61

Dividing the year in two parts:

Deaths.	Per cent.	Deaths	. Per cent.
First six months593	31.49	Second six months1,290	68.51

These figures show the great excess of mortality in the latter part of the year.

The next table shows the number and percentage of decedents from typhoid fever, in each division of age, in the aggregate mor-

tality for the	year 1892	, and als	o for the	e 37	years from	1856 to
1892, inclusiv	re:					

PERIODS.	Under 1 year.	1 and under 2.	2 and under 5.	5 and under 10	10 and under 15.	15 and under 20.	20 and under 30.	30 and under 40.	40 and under 50.	50 and under 60.	60 and under 70.	70 and under 80.	80 and under 90.	90 and over.	Total.
1892.															
Number			1	1	4	8	13	6	8	6	3	1			51
Per cent			1.96	1.96	7.85	15.69	25.49	11.76	15.69	11.76	5.88	1.96			100.00
					<u> </u>										
1856–1892.															
Number	10	17	56	116	144	294	571	280	145	125	77	38	9	1	1,883
Per cent	•53	.90	2.98	6.16	7.65	15.61	30.32	14.87	7.70	6.64	4.09	2.02	.48	.05	100.00

Of the 51 decedents from typhoid fever in 1892, 27, or 52 per cent., were between 15 and 40 years of age; of the 1,883 decedents in 37 years, 1,145, or 60.80 per cent. were between those ages.

The next table shows the number of deaths, and the aggregate and average age of the decedents in periods of five years each from 1856 to 1890; for the years 1891 and 1892; also for the 37 years, 1856-1892:

Decedents from typhoid fever; aggregate and average age.

	er of hs.	Agg	regate	Age.	Average Age.			
PERIODS.	Number of Deaths.	Years.	Mos.	Days.	Years.	Mos.	Days.	
1856-1860	141	4,358	1	16	30	10	27	
1861-1865	215	6,342	5	27	29	5	29	
1866-1870	190	5,402	8	5	28	5	6	
1871-1875	313	8,937	2	27	28	6	19	
1876-1880	216	6,030	6	1	27	11	1	
1881–1885	402	10,840	2	5	26	11	17	
1886–1890	293	8,124	7	17	27	8	22	
1891	62	1,752	5	3	28	3	5	
1892	51	1,655	1	0	32	5	13	
87 years	1,883	53,443	4	11	28	4	18	

In the report for 1883, page 80, these facts were given by single years up to that date; for 1884 and 1885, these facts may be found on page 80 of the report for 1885; for the single years from 1886 to 1890, see Report for 1890, page 78.

Heart Diseases of. The number of deaths from diseases of the heart in 1892 was 221; this was 12 more than in 1891. So far as specified these deaths were as follows: Valvular 66, heart failure 24, endocarditis 11, dropsy 10, paralysis and angina pectoris 6 each; fatty degeneration 5, hypertrophy and dilatation 4 each; rheumatism and pericarditis 2 each; myocarditis, neuralgia, enlargement, degeneration, and rupture 1 each.

Males, 110; females, 111; total, 221, or 7.49 per cent. of all.

American parentage, 97; foreign parentage, 124.

For the 37 years, 1856-1892, the number of decedents from diseases of the heart, and the particulars of sex and parentage, are given in the following table:

	Sı	EX.	PARE	TAGE.	er.	
YEARS.	Males.	Females.	American.	Foreign.	Whole number.	Percentage.
1856–1860	62	81	104	39	143	3.01
1861-1865	116	121	169	68	237	4.29
1866-1870	131	131	164	98	262	4.73
1871-1875	218	243	277	184	461	5.50
1876-1880	239	240	271	208	479	4.87
1881-1885	332	397	377	352	729	6.59
1886-1890	443	503	456	490	946	7.47
1891	101	108	82	127	209	7.99
1892	110	111	97	124	221	7-49
37 years	1,752	1,935	1,997	1,690	3,687	5.80

For particulars for single years, see reports for 1882, page 85; for 1885, page 81; and for 1890, page 79.

Heat. There were 13 deaths ascribed to heat in 1892, 11 more than in 1891. There were 3 deaths from this cause in 1890 and 1 each in 1889 and 1888. The largest number ever before recorded in one year was 7 in 1866.

Of the 13 deaths in 1892, 2 occurred in June, 9 in July and 2 in August. The sex and parentage were as follows:

Males, 11; females, 2; total, 13, or 0.44 per cent. of all. American parentage, 2; foreign parentage, 11.

Hemorrhage. The number of deaths due to this cause in 1892 was 11; 4 more than in the preceding year. Four were hemorrhage from the lungs, 4 from the umbilicus, and one each from the bowels and an injury in a case of hæmaturia. The decedents were:

Males, 7; females, 4; total, 11, or 0.37 per cent. of all, American parentage, 7; foreign parentage, 4.

Hernia was the cause of 10 deaths in 1892; 6 more than in 1891.

Males, 3; females, 7; total, 10, or 0.34 per cent. of all. American parentage, 3; foreign parentage, 7.

Homicide. Of the 5 cases of homicide, 1 was hit with a flatiron, 1 was due to pistol shot and incised wounds, 1 to pistol shot and injury to head, 1 to perforation of ileum, and 1 to stab wound of aorta.

> 1 male; 4 females. 2 of American parentage; 3 of foreign parentage.

Hydrocephalus. This was recorded as the cause of 13 deaths in 1892; the same as in 1891.

Males, 7; females, 6; total, 13, or 0.44 per cent. of all. American parentage, 5; foreign parentage, 8.

For the 37 years, 1856 to 1892, the statistics were:

Whole number, 1,025; annual average, 28.

Males, 556; females, 469; total, 1,025, or 1.01 per cent. of all.

American parentage, 523; foreign parentage, 502.

Inanition was returned as the cause of 26 deaths in 1892, 1 more than in 1891.

Males, 16; females, 10; total, 26, or 0.88 per cent. of all. American parentage, 15; foreign parentage, 11,

In the ten years, 1882 to 1892, the deaths from this cause have been:

Males, 80; females, 71; total, 151. American parentage, 76; foreign parentage, 75.

Influenza caused 59 deaths in 1892; 35 in 1891, 64 in 1890. The decedents in 1892 were:

Males, 24; females, 35; total 59, or 2.00 per cent. of all. American parentage, 35; foreign parentage, 24.

The seasonable distribution of the deaths from influenza in 1890, 1891 and 1892 was as follows:

	Jan.	Feb.	March.	April.	May.	June.	August.	Nov.	Dec.
1890	54	8	1	1	1	0	0	0	0
1891	1	0	1	1	5	4	1	0	18
1892	34	11	3	5	2	1	0	1	2

The age distribution for the three years was as follows:

		1		1	ſ	1	1	1		1		
Under 1 year	1	2-5	5-10	10-15	15-20	20-30	30-40	40-50	50-60	60-70	70-80	80-90
18902	2	0	0	4	9	12	5	6	7	11	4	2
18917	2	1	0	0	0	1	2	1	1	8	6	6
189210	1	1	1	2	0	1	2	2	7	13	9	10

Insanity. The number of deaths from this cause in 1892 was 14, or 1 more than 1891. As far as specified the varieties of insanity were as follows: Senile insanity, 8; paresis, 2.

Of the 14 deaths from insanity, only 3 were at the Butler Hospital, 11 were elsewhere in the city.

Males, 7; females, 7; total, 14, or 0.48 per cent. of all. American parentage, 8; foreign parentage, 6. In the 37 years, 1856 to 1892, the deaths from insanity numbered 396:

Males, 182; females, 214; total, 396, or 0.62 per cent, of all.

American parentage, 278; foreign parentage, 118.

Of the 396 decedents, 45.96 per cent. were males, and 54.04 per cent. females: 70.20 per cent. of American, and 29.80 per cent. of foreign parentage.

Intemperance was given as the cause of 15 deaths in 1892; 1 more than in the preceding year.

Males, 11; females, 4; total, 15, or 0.51 per cent. of all. American parentage, 1; foreign parentage, 14.

For the 37 years, 1856 to 1892:

Whole number, 390; annual average, 10.
Males, 273; females, 117; total, 390, or 0.61 per cent. of all.
American parentage, 99; foreign parentage, 291.

Of the 390 decedents, 70.00 per cent. were males; and 74.61 per cent. were of foreign parentage.

Kidneys, Diseases of. The number of deaths from diseases of the kidneys in 1892 was 122; this was 14 more than in 1891. Of the 122 deaths, 114 were specified as Bright's disease, 2 waxy kidney, and 1 each as abscess, granular kidney and pyelonephritis. The decedents were:

Males, 63; females, 59; total, 122, or 4.14 per cent. of all.

American parentage, 49; foreign parentage, 73.

For the 37 years, 1856 to 1892, the statistics were:

Whole number, 1,509; annual average, 41.
Males, 825; females, 684; total, 1,509, or 2.37 per cent. of all.
American parentage, 732; foreign parentage, 777.

Liver, Diseases of. There were in 1892, 43 deaths from diseases of the liver, which was 8 more than in the preceding year. Of these 43 cases, 14 were specified as cirrhosis, 5 as abscess, 7

hepatitis, 3 enlargement, 1 each, atrophy, congestion and amyloid degeneration. The decedents were:

Males, 18; females, 25; total, 43, or 1.46 per cent. of all. American parentage, 13; foreign parentage, 30.

For the 37 years, 1856 to 1892:

Whole number. 723; annual average, 20.
Males, 369; females, 372; total, 723, or 1.14 per cent. of all.
American parentage, 310; foreign parentage, 413.

Malformations. The number of deaths from malformation in 1892 was 14, 1 more than in 1891. They were specified as follows: Cyanosis 6, patulous foramen ovale 3, malformation of heart 4, and 1 non-contraction of ductus arteriosis. The decedents were as follows:

Males, 11; females, 3; total, 14, or 0.48 per cent. of all. American parentage, 10; foreign parentage, 4.

In the 37 years, 1856 to 1892, the decedents were:

Males, 202; females, 138; total, 340, or 0.53 per cent. of all.

American parentage, 204; foreign parentage, 136.

Marasmus was the cause given for 30 deaths in 1892; 8 more than in 1891; 22 of the decedents, or 73.33 per cent., were less than a year old.

Males, 18; females, 12; total, 30, or 1.02 per cent. of all. American parentage, 13; foreign parentage, 17.

In the 37 years, 1856 to 1892, the decedents were:

Whole number, 1,065; annual average, 29.
Males, 588; females, 477; total, 1,065, or 1.68 per cent. of all.
American parentage, 527; foreign parentage, 538.

Measles. There were 10 deaths from measles in 1892, which was 6 more than in the preceding year.

Males, 5; females, 5; total 10, or 0.34 per cent. of all.

American parentage, 3; foreign parentage, 7.

This disease has always been very irregular in its appearance in this city, as may be seen by the following table, which shows the number of deaths caused by it in each year since 1856:

8561	1865 7	1874 4	1883 8
.857	1866 9.	1875 0	1884 7
85840	1867 0	1876 0	1885 8
.859 0	1868 4	1877 2	1886
.8600	1869 6	1878 50	188790
861 6	1870 16	1879 0	1888 2
862 1	1871 1	1880 3	188914
86316	1872 7	188125	189031
864	187328	1882 4	1891 4
			189210
37 vears			422

# In the 37 years, 1856 to 1892, the decedents were:

Whole number, 422; annual average, 11.
Males, 214; females, 208; total, 422, or 0.66 per cent. of all.
American parentage, 153; foreign parentage, 269.

Meningitis, Cerebro-Spinal. The first deaths reported from this cause in Providence were in 1864; since that date the number of deaths from it each year have been as follows:

1864 2	187410	1884 6
1865 2	1875 6	1885 3
1866 1	1876 4	1886
1867 0	1877 7	1887 7
1868 1	1878 7	1888 5
1869 0	1879 7	1889 3
1870 0	1880 4	1890 3
187112	188111	1891 5
187210	188214	1892 9
187339	188315	
28 years		

#### The decedents in 1892 were:

Males, 6; females, 3; total, 9, or 0.31 per cent. of all. American parentage, 3; foreign parentage, 6.

# In 29 years, 1864 to 1892:

Whole number from known causes, 56,534.

# From cerebro-spinal meningitis:

Males, 105; females, 90; total, 195, or 0.34 per cent. of all.

American parentage, 96; foreign parentage, 99.

The number and percentage of decendents from cerebro-spinal meningitis, in each division of age, and according to sex and parentage, are given in the following table for the 29 years, 1864 to 1892:

AGES.	Under 1 year.	1 and under 2.	2 and under 5.	5 and under 10.	10 and under 15.	15 and under 20.	20 and under 30.	30 and under 40	40 and under 50	50 and under 60.	60 and under 70.	70 and nuder 80.	Males.	Females.	American.	Foreign.	Total.
No.	30	14	32	33	19	20	19	18	10	cs.	<del>,</del> 1	1	105	06	96	66	195
PER CENT.	10.26	7.18	16.41	20.00	9.74	10.26	9.74	9.23	5.13	1.03	-51	19.	53.84	46.16	49.23	22.77	100.00

Of the 195 decendents, 124, or 63.59 per cent., were less than 15 years old.

Old Age. There were 90 deaths in 1892 ascribed to this cause; 19 more than in 1891; 11 of the decedents were over 90 years of age; one of these was said to be 100 years old. The 90 decedents from old age were as follows:

Males, 27; females, 63; total, 90, or 3.05 per cent. of all. American parentage, 53; foreign parentage, 37.

# For the 37 years, 1856 to 1892:

Whole number, 2,266; annual average, 61.

Males, 750; females, 1,516: total, 2,266, or 3.56 per cent. of all.

American parentage, 1,332; foreign parentage, 934.

The decedents were 58.78 per cent. of American, and 41.22 per cent. of foreign parentage; 66.90 per cent. females, and 33.10 per cent. males.

How small a proportion of the decedents of advanced age are reported under this head will be seen by the following figures. In 1892, there were 410 decedents, or 13.83 per cent of all the decedents from known causes, who were more than 70 years old; 264 of American, and 146 of foreign parentage; 165 males and 245 females.

In the 37 years, 1856-1892, the number of decedents of 70 years and over was 7,811, or 12.29 per cent. of all the decedents from known causes; 5,201 of American and 2,610 of foreign parentage.

Paralysis. The number of deaths from this cause in 1892 was 29, which was 7 more than in 1891. Of these 29 cases, 7 were specified as hemiplegia, 4 paresis, 1 ataxic paraplegia and 1 paralysis agitans. The decedents were as follows:

Males, 9; female, 20; total, 29; or 0.98 per cent. of all. American parentage, 17; foreign parentage, 12.

In the 37 years, 1856-1892:

Whole number, 783; annual average, 21.
Males, 377; females, 406; total, 783, or 1.23 per cent. of all.
American parentage, 528; foreign parentage, 255.

The following table shows the number of deaths from paralysis and from apoplexy in each of the 37 years since 1856:

# Deaths from Paralysis and Apoplexy.

Years. Apoplexy. Paralysis. Total.	Years. Apoplexy. Paralysis. Total
18562123	1874 36 59
1857927	1875 4135
1858 318	1876
185916622	1877 50
186028	1878 42 24 60
186125	1879 56 80 80
186224	1880 48 30 78
1863	1881742710
186429	1882 8928
186538	188334118
186621	1884
1867311849	188532
1868222446	1886 99
<b>1869</b> 321648	18879927120
187052	1888 974013
1871332861	1889 822310
1872312051	18901141414
<b>1873</b> 362157	1891 892211
	1892 9829 127
Total, 37 years	

The decedents from the two causes for the 37 years, 1856 to 1892, were as follows:

Whole number, 2,579; annual average, 70.

Males, 1,231; females, 1,348; total, 2,579, or 4.06 per cent. of all.

American parentage, 1,623; foreign parentage, 956.

Of American parentage, 62.93 per cent.; of foreign parentage, 37.07 per cent.

Peritonitis. There were 14 deaths from this cause in 1892, or 13 less than in 1891.

Males, 9; females, 5; total, 14, or 0.48 per cent. of all. American parentage, 7; foreign parentage, 7.

## In the 37 years, 1856 to 1892:

Whole number, 496; annual average, 13.

Males, 209; females, 287; total, 496, or 0.78 per cent. of all.

American parentage, 244; foreign parentage, 252.

Phthisis, or Consumption. There were 333 deaths from phthisis in 1892, or 11.29 per cent. of all deaths from known causes; the number was 11 less, and the rate 1.86 per cent. less than in 1891.

The population of Providence for the year 1892 was estimated at 138,000; the 333 deaths from phthisis therefore average one in each 414.4 of the inhabitants, or 241.3 in each 100,000. In 1891 the rate was one in 3,924, or 254.8 in each 100,000.

During the 15 years, 1840 to 1854, the average annual rate of mortality from phthisis was one death in each 209.9 inhabitants, or 476 to each 100,000. During the 37 years, 1856 to 1892, the annual average rate for phthisis was one death to each 328.6 inhabitants, or 304.4 to each 100,000.

## In 1892 the decedents from phthisis were as follows:

American parentageMales,		
Foreign parentage Males,	113Females,	127Total, 240
TotalsMales,	165Females,	168Total, 333

# The statistics for the 37 years, 1856-1892, are:

Total deaths from known causes. 63,571; from phthisis, 9,852, or 15.50 per cent.

Annual average number, 266.

American parentage.......Males, 1,791.......Females, 2,177.......Total, 3,968

Foreign parentage.......Males, 2,839........Females, 3,045........Total, 5,884

Totals ........Males, 4,630.......Females, 5,222......Total, 9,852

Males 47.00.....Females, 53.00 in each 100.

American parentage, 40.28......Foreign parentage, 59.72 in each 100.

The table on the next two pages gives the particulars of sex, parentage, season and locality, in regard to the decedents from phthisis in the five years, 1856 to 1860; the five years, 1861 to 1865; the five years, 1866 to 1870; the five years, 1871 to 1875; and in each year since 1875:

DEATHS FROM PHTHISIS IN PROVIDENCE, THIRTY-SEVEN YEARS, 1856-1892.

Sex; parentage; season; age; whole number and ratio to known causes.

Totals.	9,852 15.50	4,630 5,222	3,368	931 866 874 8836 8836 8836 765 765 816 816 835 835
1892.	333	165	93	######################################
1891.	344	188	98	£&\$\$\$\$\$\$\$\$\$\$\$\$\$\$
1890.	385 13.46	205	271	32423333333333
1889.	309	155	104	%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
1888.	359 13.80	183 176	98	######################################
1887.	314	141 173	98	88888888888
1886.	368 15.66	179	253	58888888888888888888888888888888888888
1885.	347	176	124	448888888888
1884.	344	159	231	888 4888888888 888 4888888888
1883.	364 15.68	166	107	828644888828828
1882.	351	166	113	48888888888888
1881.	344	157	228	<u> </u>
1880.	322	154	2111	828222888888
1879.	293	141	116	828884488888
1878.	305	135	107	22222222222222222222222222222222222222
1877.	294	127	116	88444
1876.	284	119	120	### ### ### ### ### ### ### ### ### ##
1871 to 1875.	1,235	568	553 682	110 101 101 101 102 88 88 109 109 109 109 109 109 109 109 109 109
1865 to 1870.	1,052	490	532	88888888888888888888888888888888888888
1861 to 1865.	975	477	495	F1888688F18888
1856 to 1860	930	379 551	525 405	88 88 88 88 88 88 88 88
PHTHISIS.	Whole number	SEX: Males Females	PARENTAGE: American Foreign	SEASON: January February March April May June July September October November December

DEATHS FROM PHTHISIS IN PROVIDENCE, THIRTY-SEVEN YEARS, 1856-1892.—CONCLUDED.

Sex; parentage; season; age; whole number and ratio to known causes.

			*
Totals.	2.592 2,513 2,282 2.465	3,466 3,158 3,228	131 100 100 92 92 103 103 8712 8712 872 873 873 873 873 873 873 873 873 873 873
1892.	28.00 7.27 7.25	1119	481-85-838-881
1891.	2818	121 107 116	800 00 00 00 00 00 00 00 00 00 00 00 00
1890.	114 94 88 93	130	5-66.85.888.88.25.7.4
1889.	77.8 20.0 7.1 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0	103	-0000000000000000000000000000000000000
1888.	8888	121 118 110	701 4 80 80 80 4 1 37 57 57 50 60 60 60 60 60 60 60 60 60 60 60 60 60
1887.	200 00 00 00 00 00 00 00 00 00 00 00 00	117	4103458865885574
1886.	104 1114 877 63	149 132 87	<u></u>
1885.	25 2 2 2	1125	20014811 100014811888881001
1884.	103 173 88	1110	1128 8 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
1883.	1068	119	25.00 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
1882.	102 24 75 80	1115	9384 4 8 8 6 6 8 8 6 6 8 8 8 6 8 8 8 8 8 8
1881.	2828	125 123 96	44005-1555-2851
1880.	8288	90 90 117	444-8887599 66-6-6-6-6-6-6-6-6-6-6-6-6-6-6-6-6-6-
1879.	42668	112 89 92	1000408888888551
1878.	22826	91 103 111	L4L4@150088331
1877.	1382	103 100 100	4-84r070304482F
1876.	35.25	86 107 91	44-600 B B B B B B B
1871 to 1875.	327 317 313 313	444 382 409	111112120131111111111111111111111111111
1866 to 1870.	277 270 249 262	365 343 344 344	52552888825525 525528888855
1861 to 1865.	242 227 239 267	328 296 351	44102202222222 441022022222222222
1856 to 1860	265 190 265 265	328 331	200 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
PHTHISIS.	First quarter Second quarter Third quarter Fourth quarter	First 4 months Second 4 months Third 4 months	AGE: Under I year. 2 and under 5. 5 and under 5. 15 and and 5. 15 and and 5. 15 and and 6. 16 and and 6. 16 and and 6. 17 and and 6. 18

The following figures give the number and percentage, according to parentage, of the decedents from phthisis, in periods of five years, from 1856 to 1890, in 1891, 1892, and in the whole period of 37 years:

	American	parentage.	Foreign p	arentage.	Total from
	Number.	Per cent.	Number.	Per cent.	Phthisis
56 to 1860	525	56.45	405	43.55	95
61 to 1865	495	50.77	480	49.23	9
66 to 1870	532	50.57	520	49.43	1,0
71 to 1875	553	44.78	682	55.22	
76 to 1880	570	38.05	928	61.95	1,4
81 to 1885	573	32.74	1,177	67.26	
86 to 1890	529	30.49	1,206	69.51	
1891	98	28.49	246	71.51	3
7.000	00	27.93	240	70.07	3:

This table shows what a very great change has taken place in the mortality from phthisis in Providence in the last thirty-six years, in regard to parentage; the proportions of American and foreign parentage among the decedents being quite the reverse now from what they were in 1856. This change was very fully illustrated and commented upon, under the heading consumption, in the report for 1881.

Pneumonia. There were 304 deaths from pneumonia in 1892, and 11 from congestion of the lungs; total, 315, which was 64 more than in the preceding year. These deaths were as follows in the different seasons of the year:

Number.	Per cent.	Number. Per cent.
First quarter158	50.16	Third quarter17 5.40
Second quarter 71	22.54	Fourth quarter6921.90

#### The decedents were:

Males, 160; females, 155; total, 315, or 10.68 per cent. of all American parentage, 103; foreign parentage, 212.

In the 37 years, 1856 to 1892, the decedents from pneumonia were:

Whole number, 5,131; annual average, 139.

Males, 2,624; females, 2,507; total, 5,131, or 8.07 per cent. of all.

American parentage, 2,346; foreign parentage, 2,785.

The following table gives the statistics concerning the decedents from pneumonia and congestion of the lungs, in the five years, 1856–1860; the five years, 1861–1865; the five years, 1866–1870; and in each year since 1870. See the report for 1883, pages 89 to 93 inclusive, for fuller particulars and comments:

PNEUMONIA AND CONGESTION OF THE LUNGS IN PROVIDENCE, THIRTY-SEVEN YEARS, 1856-1892.

Sex; parentage; season; age; whole number and ratio to known causes.

	Totals.	5,131	2,624 2,507	2,346	74 1178 1178 1178 1178 1178 1178 1178 11
	1892.	315	155	103	%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
Ì	1891.	251 9.59	127	101	4921 912 912 912 912 912 912 912 912 912
	1890.	244 8.53	130	88 156	28 E 8 E E E E E E E E E E E E E E E E E
	1889.	204 8.19	100	121	00847-c0000000
	1888.	221 8.48	115	131	884880446817
	1887.	224 8.60	123	88 136	88488850005342
	1886.	210 8.93	103	93	23.23.23.23.23.23.23.23.23.23.23.23.23.2
	1885.	227 10.51	113	124	<u> </u>
	1884.	172	97.8	888	
	1883.	204 8.79	109	89	271888888 000 000 000 000
	1882.	176	28	99	71 88 86 86 86 86 86 86 86 86 86 86 86 86
	1881.	174 8.17	20 00 10 00	88	285245150005555
	1880. 1881.	193 9.36	98	107	71 68 88 84 80 10 10 10 10 10 10 10 10 10 10 10 10 10
	1879.	159 7 88	33.75	7.3	88888888888888888888888888888888888888
	1876. 1877. 1878. 1879.	140	11.68	5.5	11121356641112
	1877.	5.07	44.	51	2111000 4 8 H 2 0 0 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
	1876.	163	79	102	21 10 10 10 10 10 10 10 10 10 10 10 10 10
	1871 to 1875.	653	334 319	334 319	2382123267
	1866 to 1870.	7.24	214 193	205	25 25 26 26 27 27 27 27 27 27 27 27 27 27 27 27 27
	1861 to 1865.	355	190	193	946888888888888888888888888888888888888
The state of the s	1856 to 1860.	341	192 149	184 157	23.01.00.00.00.00.00.00.00.00.00.00.00.00.
	PREUMONIA AND CONGRESTION OF THE LUNGS.	Whole number	SEX: Males Females	PARENTAGE: American Foreign	SEASON: January February March April May June June June September October November

PREUMONIA AND CONGESTION OF THE LUNGS IN PROVIDENCE, THIRTY-SEVEN YEARS, 1856-1892.—CONCLUDED.

causes.
known
to
and ratio
and
number
whole
age;
season;
parentage;
Sex;

Totals.	2507 116 116 116 128 128 138 138 138	2,027 1,394 445 1,265 2,670 1,023 1,438
1892.	24128282828211 28888888211	158 17 17 17 69 50 50 76
1891.	825484841488894	76 103 103 103 107
1890.	25.50 25.50	108 552 559 131 664 441
1889.	62 0114 : 001 002412821	97 171 121 121 888 455
1888.	71147	112 555 14 14 30 80 84
1887.	8118comm1888884	102 813 82 83 83 83 83 83 83 83 83 83 83 83 83 83
1886.	£021-870888848800L	-
1885.	8274001108888418	88 828 116 116 126 41 41 41 41
1884.	9m14-m4mees811s	
1883.		_
1882.	1082282113020011	651 61 61 64 64 64 65 65 65 65 65 65 65 65 65 65 65 65 65
1881.	8881-81-4488800	014 010 04 488 488 488
1880.	%;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;	2544 244 244 254 254
1879	120 120 0 0 120 0 1	48.00 88.00 44 48.00 88.00 44
1878.		4 8 3 3 1 3 9 8 9 8 9 8 9 9 9 9 9 9 9 9 9 9 9 9 9
1877.	1221 1230 1230 1230 1330 1330 1330 1330	885 498 898
1876.	24 C L C L C L C L C C L C C L C C L C C L C C L C C L C C L C C L C C C L C	4488 4988 44888
1871 to 1875.	44641148277488671°	253 174 68 158 158 136 136 187
1866 to 1870.	29820428814881	1126 1128 46 93 218 73 110
1861 th 1865.	%E824%%%%%	136 99 98 171 171
1856 to 1860.	845 E. C. C. C. S.	123 977 88 157 101
PNEUMONIA AND CONGESTION OF THE LUNGS.	AGE:  2	First quarter Second quarter. Third quarter Fourth quarter First 4 months Second 4 months.

Full statistics have already been given as to the mortality from bronchitis. That disease is so closely allied to pneumonia that, while the statistics of each are kept quite distinct, it is well also to combine them, as forming one class of diseases. In 1892 the decedents from the two causes, bronchitis and pneumonia, were:

Males, 224; females, 239; total, 463, or 15.69 per cent. of all. American parentage, 161; foreign parentage, 302.

For the 37 years, 1856 to 1892, the statistics for the two causes are:

Whole number, 6,655; annual average, 180.

Males, 3,316; females, 3,339; total, 6,655, or 10.47 per cent. of all.

American parentage, 3,007; foreign parentage, 3,648.

Rheumatism. There were 13 deaths ascribed to this cause in 1892; 3 less than in 1891.

Males, 7; females, 6; total, 13, or 0.44 per cent. of all. American parentage, 6; foreign parentage, 7.

In the 37 years, 1856–1892:

Males, 110; females, 146; total, 256, or 0.40 per cent. of all.

American parentage, 109; foreign parentage, 147.

Scarlet Fever. There were 37 deaths from scarlet fever in 1892; this was 20 more than in 1891. The number and proportion in each quarter of the year were as follows:

Number. Per cent.	Number. Per cent.
First quarter1540.54	Third quarter38.11
Second quarter 8 21.62	Fourth quarter1129.73

Males, 22; females, 15; total, 37, or 1.26 per cent. of all. American parentage, 14; foreign parentage, 23. For the 38 years, 1855 to 1892, the decedents were:

Whole number, 2,568; annual average, 69.
Males, 1,246; females, 1,322; total, 2,568, or 3.98 per cent. of all.
American parentage, 1,041; foreign parentage, 1,527.

The following table shows the particulars of sex, parentage, season and locality, and the ratio to known causes, of the decedents from scarlet fever for the year 1855, for the five years, 1856–1860; the five years, 1861–1865; the five years, 1866–1870; and for each year since 1870:

DEATHS FROM SCARLET FEVER IN PROVIDENCE, THIRTY-EIGHT YEARS, 1855-1892.

Totals.	3.98	1,246	1,041	232 230 230 230 230 231 231 231 231 231 231 231 231 231 231	362 849 109 260 259 259 272 272 272 301
1892.	1.26	15	48	41001410101 : 45-	. # 4 . D W 4 7 D W 7 D
1891.	0.65	ж G	10	_ :vos : : : : : : : : : :-	
	0.28	44	0.00		
1889. 1890.	1.01	133	20	4 : wow ws 4 : : H	-22::::52
1888.	3.08	39	43	810000000000000000000000000000000000000	2002222003
1887.	153	87	8621	884197080774888	<u></u>
1886.	1.28	110	22.9	400:110004100410	: 20 HH4H888H
1885.	38	25.	12 26	70004 et-eten-co	
1884.	2.56	8.22	83	9 100 9400 00 1110	F-8000484444
1883.	$\frac{16}{0.69}$	<u>}-</u> 5	10	cs : cs cs : : : cs 44	
1882.	0.81	12	11	0100mm0001 01 mm	9:-: %-9
	2.11	18	17.	700045-0044H H05H	: :400000470
1880. 1881	243	122	155	0.0000000000000000000000000000000000000	11-37-231-822-9
1879.	252	135	101	350000 4 ro 0 0 8 4 E	82.8814018848
1878.	57.89	888	17.	ಪಪಪಲಪಾಧವಾಣಪಡೆಯ	, , , , , , , , , , , , , , , , , , ,
1877.	12 0.62	4100	2000	w : 03H : : 403	
1876.	35	15	14 21	400 410	4 . : === 0.00 5: 44
	4.91	53	653	00000000400F0	ಬ್ಯಾಂದ – ಇದೆ ಅಸ್ಥ
1874. 1875.	268 13.66	140	191	%24242423	14.00 to 22
1873.	132	7.0	7.58	%araciicare	38 27 28 20 17 28 28
1872.	1.30	118	11.8		ग्राच्या । ग्राच्या
1871.	1.69	20∞	9 31	©₽-4H65H · · · · · · · · · · · · · · · · · · ·	H 63 4 H 10 65 65 10 10
1866 to 1870.	236	106	98	222256-7227882	22 22 22 23 25 25 25
1861 to 1865.	324 5.86	154	132	7.E888888111884	42 188 244 377 111
1856 to 1860.	310	153	172	8684488881-xeee	25 10 10 10 10 10 10 10 10 10 10 10 10 10
1855.	4.37	38	13	:	91987
SCARLET FEVER.	hole number	Males	American	ASON: January January Rebruary March April April April August Seeptember October December	Vard II Vard II Vard VII VIII VIII VIII VIII VIII

The ward totals in this table do not include the figures after 1887, as those are reckoned by the new division of wards; and all before that are for the old wards of the city.

Scrofula was returned as the cause of 1 death in 1892; 4 less than in 1891.

Males, 1; females, 0; total 1, or 0.03 per cent. of all.

American parentage, 1; foreign parentage, 0.

Small-Pox. There were 3 deaths from this disease in 1892, the first since 1883. The decedents were:

Males, 2; female, 1, total 3, or 0.10 per cent. of all. American parentage, 0; foreign parentage, 3.

The following are the statistics for 37 years from 1856-1892:

Males, 61; females, 22; total, 83, or 0.13 per cent. of all. American parentage, 49; foreign parentage, 34.

The following table shows the particulars for each year:

#### SMALL-Pox.

	8	SEX.	PARE	NTAGE.	Whole	
Year.	Males.	Females.	American.	Foreign.	Number.	Percentage
1856	7	2	5	4	9	0.8
1857						
1858						
1859	2	3	5		5	
1860	4	1	3	· · · · · · · · · 2 · · · ·	5	
1861						
1862	2	2	1		4	0.4
1863		/				
1864	7		3	4	7	
		2				
1866			******			
1867	1		1		1 .	0.1
1868	1		1		1	0.0
1869	3		3			
1870		2		2		
1871		3		2	3	
1872	9	4	9	4	13	
		2				
1874						
1875	1			1	1	0 . 0
1876						
1877						
1878						
1879						
1880						
1881						
1882						
	1		1			
1884						
1885						
1886						
1887						
1888						
1889						
1890						
1891						
				3		
1094	2	1				

Evidences of successful vaccination are required of all children entering the public schools of Providence.

Spinal Cord, Diseases of. There were 10 deaths in 1892, specified as follows: locomotor ataxia, 4; tabes dorsalis, 4; acute ascending paralysis, 1. The decedents were, males 8; females 2; total 10 or 0.34 per cent. of all. American parentage 3; foreign parentage 7.

Spine, Diseases of. There were 4 deaths in 1892 from diseases of the spine; 2 Potts disease, and 1 each, necrosis of coccyx and caries of cervical vertebrae.

Males, 1; females, 3; total, 4, or 0.14 per cent. of all. American parentage, 1; foreign parentage, 3.

Stomach, Diseases of. The number of deaths from diseases of the stomach in 1892 was 26, of which 20 were called "gastritis." The decedents were:

Males, 14; females, 12; total, 26, or 0.88 per cent. of all. American parentage, 6; foreign parentage, 20.

Suicide. There were 8 deaths by suicide in 1892, 16 less than in 1891. Two were by shooting with pistol and one each by suffocation with illuminating gas, poisoning with laudanum, by opium, by cutting the throat, by strangulation, and by drowning.

Males, 7; females, 1; total, 8, or 0.27 per cent. of all. American parentage, 3; foreign parentage, 5.

In the 37 years, 1856–1892, the decedents by suicide were:

Males, 207; females, 69; total, 276, or 0.43 per cent. of all.

American parentage, 142; foreign parentage, 134.

Of the 276 decedents, 75.00 per cent. were males; and 51.45 per cent. were of American Parentage.

Surgical Operation. There were 6 deaths in 1892 from surgical operation, 5 more than in 1891. Of these there was one each from operation for harelip, from laparatomy, ovairotomy, suprapubic cystotomy, operation for cancer of the uterus and for cyst of the broad ligament.

The decedents were:

Males, 2; Females, 4; total 6, or 0.20 per cent. of all. American parentage, 4; foreign parentage, 2.

Teething was given as the cause of 8 deaths in 1892, or 14 less than in 1891.

Males, 5; Females, 3; total, 8, or 0.27 per cent. of all. American parentage, 3; foreign parentage, 5. Tubercolosis was the cause of 37 deaths in 1892, 11 more than in 1891.

Males, 17; females, 20; total 37, or 1.25 per cent. of all. American parentage, 20; foreign parentage, 17.

The name "tuberculosis" was first used in these reports in 1867; in the 26 years, 1867 to 1892, the deaths from it have been as follows:

All deaths from known causes in 26 years, 52,279.

From tuberculosis, 469; annual average, 18.

Males, 225; females, 244; total, 469, or 0.90 per cent, of all.

American parentage, 274; foreign parentage, 195.

If under tuberculosis should also be included tubercular meningitis, phthisis and hydrocephalus, it is probable that some few cases of non-tuberculous disease might be included, but as a certain number of cases of true tuberculosis are undoubtly reported as adenitis, disease of bone, peritonitis, etc., it is probable that this total would be below rather than above the actual number of deaths from tubercular disease. The total of these tubercular diseases for 1892 was 412, or 13.97 per cent. of deaths from known causes.

Tubercular Meningitis. The number of deaths from this cause in 1892 was 29.

Males, 17; females, 12; total, 29, or 0.98 per cent. of all. American parentage, 16; foreign parentage, 13.

Tumors. The number of deaths from tumors in 1892 was 6, or 10 less than in 1891. There were 4 ovarian; 1 abdominal; 1 fibro-sarcoma uteri. The decedents were:

Males, 0; females, 6; total, 6, or 0.20 per cent. of all. American parentage, 3; foreign parentage, 3.

Whooping Cough. There were 8 deaths from whooping cough in 1892, 45 less than in the preceding year. The decedents were:

Males, 5; females, 3; total, 8, or 0.24 per cent. of all. American parentage, 4; foreign parentage, 4. In the 37 years, 1856 to 1892, there were:

Whole number, 731; annual average, 20.
Males, 334; females, 397; total, 731, or 1.15 per cent. of all.
American parentage, 321; foreign parentage, 410.

Unknown. There were 14 deaths in 1892, for which no cause could be given. This was the same as in 1891; it was 0.47 per cent. of the 2,964 deaths recorded for the year. The decedents were:

Males, 3; females, 11; total, 14, or 0.47 per cent. of all. American parentage, 2; foreign parentage, 12.

For the 37 years, 1856 to 1892, the statistics are:

Whole number, 676; annual average, 18.
Males, 344; females, 332; total, 676, or 1.05 per cent. of all.
American parentage, 239; foreign parentage, 437.

Of the 676 decedents, 64.64 per cent. were of foreign parentage; and 347, or 51.33 per cent. were children under 1 year of age.

# STILL-BORN.

There were 190 still-born children in Providence in 1892; this was 17 more than in 1891, and was 4.81 per cent. of the 3,944 children born in 1892, or one in 20.7.

Sex. Males, 113; females, 77.

Color. White, 185; colored, 5.

Locality. First ward, 13; second, 12; third, 36; fourth, 9; fifth, 22; sixth, 14; seventh, 7; eighth, 19; ninth, 24; tenth, 34.

Season. January, 11; February, 13; March, 16; April 12; May, 20; June 13; July, 22; August, 14; September, 14; October, 19; November, 20; December, 16.

The number and proportion in each quarter were:

	Number.	Per cent.	Number. Per cent.	
First quarter	40	21.05	Third quarter5026.32	
Second quarter	45	23.68	Fourth quarter5528.95	

Parentage. The following table shows the parentage of the 190 still-born children in 1892; the first letter representing the birth-place of the father, the second that of the mother.

A.—American; I.—Irish; E.—English; S.—Scotch; G.—German; B. A.—British American; F. C.—French Canadian; P.—Portuguese; Sw.—Swedish; It.—Italian; R. Russian; Aus—Austrian; Syr.—Syrian; Pol.—Polish; N.—Norwegian; Rou.—Roumanian.

# Parentage.

A. and A59	Pol. and Pol 1	I. and E 1
I. and I44	Syr. and Syr 1	S. and E 1
E. and E 8	I. and A 8	B. A. and E 1
S. and S 1	E. and A 3	A. and S 2
G. and G 4	B. A. and A 1	Swiss and S 2
B. A. and B. A 3	N. and A 1	R. and G 1
F. C. and F. C 2	S. Hel. and A 1	A. and B. A 3
Sw. and Sw 2	W. Ind. and A 1	A. and Maltese 1
It. and It 6	A. and J10	R. and Rou 1
P. and P 3	E. and I 5	comments
R. and R 5	A. and E 8	Total190

The number and proportion of children borne by mothers of different nativities in 1892, were as follows:

	Number of children.	Per cent.
American mothers		40.87
Irish mothers	871	
Other foreign mothers	,	
Unknown	7	
	- <del></del>	
Total		100.00

The number and percentage of children of American parentage was less in 1892 than in 1891. Of both the other classes, the number and percentage were more in 1892.

The next table shows the same facts concerning the 190 still-born children in 1892.

	Still-born children.	Per cent.
American mothers	74	38.95
Irish mothers	59	31.05
Other foreign mothers	57	30.00
Total		100.00

The proportion of still-born children to the whole number of children borne by mothers of each class in 1892, is shown by the following table:

American mothers		children.	Percentage of still-born.	
Irish mothers Other foreign mothers	871	59	6.77	14.8
Unknown				-

In all classes the proportion of still-born children to the whole number of children born was more in 1892 than in the preceding year.

In the 37 years, 1856 to 1892, the whole number of children born was 86,985; of still-born, 4,910, or 5.64 per cent., or 1 in 17.7.

Of the 4,910 still-born children, 4,617 were white and 293 colored; 2,830, or 57.64 per cent. were males, and 2,079, or 42.34 per cent. were females; in 1891, the sex of one was not stated.

The following statement shows the whole number of children borne by mothers of different nativities during the 37 years 1856–1892, inclusive; also the number of still-born children to mothers of each class; and the percentage of the latter to the whole number of children.

	Whole number		Percentage	
			of still-born.	
American mothers		2,191	5.31	18.8
Irish mothers,		1,882	6.66	15.0
English, Scotch or Welsh m	others. 7,805	421	5.39	18.5
German mothers	1,816	66	3.63	27.5
British-American mothers	3,766	176	4.67	21.4
Other foreign mothers	4,069	168	4.13	. 24.2
Unknown	16	6		
Total	86,985	4,910	5.64	17.7

Reckoned by the fathers' birth-places, 2,205 or 44.91 per cent. of the 4,910 still-born children were American; 1,784, or 36.33 per cent. were Irish; and 907, or 18.47 per cent. were children of other foreign fathers; of 14, or 0.29 per cent. the parentage was not stated.

# GENERAL SUMMARY.

The following recapitulatory table gives the general results of registration in Providence during the 38 years 1855 to 1892 inclusive:

TABLE XXI. BIRTHS, MARRIAGES AND DEATHS.

Deaths in each 1,000,	20.74	21.86	18.70	20.56	17.94	19.76	20.74	17.50	22.90	23.81	22.18	18.60	16.89
noitslugod 10 ai atsəb əno	48.22	45.75	53.46	48.63	55.72	50.61	48.21	57.15	43.67	42.01	45.08	53.77	59.19
реатрв.	166	1,065	925	1,017	899	1,001	1,051	914	1,214	1,281	1,211	1,036	096
Persons mar- ried in each 1,000,1	26.62	26.93	24.83	22.65	24.59	24.99	21.59	20.94	23.01	27.47	25.97	29.15	30.09
Of population one person married in	37.56	37.14	40.27	44.15	40.66	40.02	46.31	47.75	43.46	36.40	38.50	34.30	33.23
Marriages.	989	929	614	260	919	633	547	547	610	739	50%	812	855
Births in each 1,000.	33.49	34.38	34.13	34.87	31.80	32.53	34.05	29.27	24.65	24.97	23.78	29.29	28.58
of population of direction	29.86	29.09	29.30	28.68	31.45	30.74	29.37	34.16	40.56	40.04	42.06	34.13	34.99
Births.	1,000	1,675	1,688	1,724	1,593	1,648	1,725	1,529	1,307	1,344	1,298	1,632	1,624
Population.	147,785	48,727	49,457	49,457	50,100	*50,666	20,666	52,238	53,024	53,810	154,595	55,709	56,824
YEAR.	855	850	857	858			861	862		708	865	866	

# GENERAL SUMMARY.—CONTINUED.

The following recapitulatory table gives the general results of registration in Providence during the 38 years 1855 to 1892 inclusive:

TABLE XXI. BIRTHS, MARRIAGES AND DEATHS.

Deaths in each 1,000,	17.31	18.88	18.33	17.41	21.34	21.98	20.60	19.03	18.37	19.19	19.44	19.56	19.84
nottsluqoq tO ni disəb əno	57.78	52.96	54.55	57.42	46.86	45.51	48.54	52.57	54.43	52.12	51.43	51.12	50.41
<b>.</b> Веаthа.	1,110	1,256	1,263	1,254	1,603	1,719	1,987	1,915	1,865	1,938	1,989	2,026	2,080
Persons mar- ried in each 1,000,	27.20	96.12	27.83	26.23	28.88	29.40	24.38	21.44	18.70	18.81	19.86	89.02	23.53
nottaluqoq 10 nostaq ano ni bairram	36.77	35.76	35.93	38.13	34.63	34.01	41.01	46.65	53.48	52.99	50.34	48.35	42.49
.яэувіттвМ	872	930	959	943	1,084	1,150	1,176	1,079	949	953	1,016	1,071	1,234
Births in each 1,000,1	60.62	27.34	27.41	27.22	29.37	27.30	28-83	26.46	25.92	25.18	25.68	24.35	25.05
of hypopulation of of direction	34.37	36.57	36.48	36.74	34.05	36.76	34.69	87.79	38.58	39.72	38.94	41.07	39.92
Births.	1,866	1,819	1,889	1,960	2,206	2,128	2,780	2,664	2,631	2,543	2,627	2,522	2,627
Population,	64,138	66,522	*68,904	72,011	75,118	78,225	196,452	†100,675	101,511	101,000	\$102,286	103,572	*104,857
YEAR	898	869,	0.28		8.28	(8.5)		SWZ		Man	1878		1880

GENERAL SUMMARY.—CONCLUBED.

The following recapitulatory table gives the general results of registration in Providence during the 38 years 1855 to 1892 inclusive:

TABLE XXI. BIRTHS, MARRIAGES AND DEATHS

YEAR.	Population.	Births.	nottslugog 10 of dirid ano	Births in each	. казавіттыМ	noitsluqoq 10 noereq eno ni betrism	Persons mar- ried in each 1,000,	Deaths.	noitaluqoq 10 ni diasab sno	Deaths in each
1881	109,617	2,803	39.10	25.58	1,202	45.60	21.93	2,145	51.10	19.56
1882	114,377	2,788	41.03	24.38	1,172	48.80	20.49	2,242	51.01	19.60
1883	#117,018	2,916	40.13	24.92	1,177	49.71	20.12	2,328	50.27	19.89
1884	117,544	2,956	39.76	25.15	1,144	51.37	19.47	2,233	52.64	19.00
1885	+118,070	2,936	40.21	24.87	1,091	54.11	18.48	2,164	54.56	18.33
1886	120,000	2,960	40.54	24.67	1,249	48.04	20.82	2,355	50.96	19.62
1887	121,500	2,963	41.01	24.38	1,336	45.47	21.99	2,612	46.52	21.50
1888	123,000	3,131	39.28	25.46	1,349	45.59	21.93	2,608	47.16	21.20
1889	127,000	3,171	40.05	24.97	1,367	46.45	21.53	2,510	50.60	19.76
1890	*132,146	3,417	38.67	25.86	1,430	46.20	21.64	2,877	45.93	21.77
1891	135,000	3,951	34.17	29.27	1,488	45.36	22.04	2,630	51.33	19.48
1892	138,000	3,944	34.99	28.58	1,592	43.34	23.72	2,964	46.56	21.48
38 years, average	86,358	2,331	37.05	26.99	186	43.75	22.86	1,717	50.29	19.88
*	co		† State	State Census.		+-	City Conging	2112	,	

#### BIRTHS AND DEATHS BY PARENTAGE.

The following recapitulatory table shows the number of births and deaths by parentage, in Providence, in each of the 37 years, from 1856 to 1892, inclusive. In the three years 1863, 1864 and 1865, there was an excess of deaths over the births, in the population of American parentage:

TABLE XXII.

	AMERIC	CAN PARI	ENTAGE.	FOREIGN PARENTAGE.				
YEAR.	Deaths.	Births.	Excess of Births.	Deaths.	Births.	Excess of Births.		
1856	541	759	218	524	916	392		
1857	523	701	178	402	987	585		
1858	512	771	259	505	953	448		
1859	479	659	180	420	934	514		
1860	512	731	219	489	917	428		
1861	538	740	202	513	985	472		
1862	460	663	203	454	866	412		
1863	597	528	-69	617	779	162		
1864	649	579	-70	632	765	133		
1865	631	546	-85	580	752	172		
1.866	513	726	213	523	906	383		
1867	528	725	197	432	899	467		
1868	596	882	286	514	984	470		
1869	627	926	299	629	893	264		
1870	642	890	248	621	999	378		
1871	643	980	337	611	980	369		
1872	832	1,035	203	771	1,171	400		
1878	862	1,003	141	857	1,125	268		
1874	885	1,217	332	1,102	1,563	461		
1875	914	1,180	266	1.001	1,484	483		
1876	875	1,169	294	990	1,462	472		
1877	908	1,238	330	1,030	1,305	275		
1878	889	1,215	326	1,100	1,412	312		

# BIRTHS AND DEATHS BY PARENTAGE.

# TABLE XXII.—CONCLUDED.

	AMERIC	CAN PARE	ENTAGE.	FOREIGN PARENTAGE.				
YEAR.	Deaths.	Births.	Excess of Births.	Deaths.	Births.	Excess of Births.		
1879	941	1,228	287	1,085	1,294	209		
1880	920	1,252	332	1,160	1.375	215		
1881	999	1,365	366	1,146	1,438	292		
1882	965	1,370	405	1,277	1,418	141		
1883	1,047	1,457	410	1,281	1,459	178		
1884	992	1,452	460	1,241	1,504	263		
1885	1,012	1,447	435	1,152	1,489	337		
1886	1,029	1,469	440	1,326	1,491	165		
1887	1,082	1,391	309	1,530	1,572	42		
1888	1,101	1,544	443	1,507	1,587	80		
1889	1,073	1,441	368	1.437	1,730	293		
1890	1,195	1,513	318	1,682	1,904	222		
1891	1,083	1,698	615	1,547	2,178	631		
1892	1,179	1,601	422	1,783	2.251	468		
Thirty-seven years	29,774	40,091	10,317	34,471	46,727	12,256		

Respectfully submitted,

CHARLES V. CHAPIN,

City Registrar.



